

ty GENUINE 240 County Road

Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: BstZ17I-HF®
Catalog Number: R3594L
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of λ DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10107916
Expiration Date: 01/2023
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl , 50 mM KCl , 1 mM DTT , 0.1 mM EDTA , 200 µg/ml BSA

, 50 % Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-R3594S/L v2.0

BstZ17I-HF® Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R3594LVIAL	BstZ17I-HF®	10097044	Pass	
B7024AVIAL	Gel Loading Dye, Purple (6X)	10093123	Pass	
B6004SVIAL	rCutSmart™ Buffer	10107577	Pass	

Assay Name/Specification	Lot # 10107916
Endonuclease Activity (Nicking) A 50 µl reaction in CutSmart® Buffer containing 1 µg of supercoiled PhiX174 DNA and	Pass
a minimum of 20 units of BstZ17I-HF incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and a minimum of 100 units of BstZ17I-HF incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
Protein Purity Assay (SDS-PAGE) BstZ17I-HF is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Ligation and Recutting (Terminal Integrity) After a 20-fold over-digestion of Lambda DNA with BstZ17I-HF, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BstZ17I-HF.	Pass



R3594L / Lot: 10107916

Page 1 of 2

Assay Name/Specification	Lot # 10107916
Functional Testing (15 minute Digest) A 50 µl reaction in CutSmart® Buffer containing 1 µg of Lambda DNA and 1 µl of BstZ17I-HF incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in CutSmart® Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 100 units of BstZ17I-HF incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghaa Zhang Production Scientist

08 Jun 2021

Michael Tonello

Packaging Quality Control Inspector

08 Jun 2021



R3594L / Lot: 10107916

Page 2 of 2