

## New England Biolabs Certificate of Analysis

**Product Name:** Bsal-HF®v2  
**Catalog Number:** R3733S  
**Concentration:** 20,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of pXba DNA in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10177238  
**Expiration Date:** 08/2024  
**Storage Temperature:** -20°C  
**Storage Conditions:** 20mM Tris-HCl, 300mM NaCl, 0.1mM TCEP, 200 µg/ml rAlbumin, 50% Glycerol, (pH 9.0 @ 25°C)  
**Specification Version:** PS-R3733S/L v2.0

Bsal-HF®v2 Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R3733SVIAL	Bsal-HF®v2	10156743	Pass
B7024AVIAL	Gel Loading Dye, Purple (6X)	10173660	Pass
B6004SVIAL	rCutSmart™ Buffer	10175290	Pass

Assay Name/Specification	Lot # 10177238
<p><b>qPCR DNA Contamination (E. coli Genomic)</b>            A minimum of 20 units of Bsal-HF®v2 is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass
<p><b>RNase Activity (Extended Digestion)</b>            A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 20 units of Bsal-HF®v2 is incubated at 37°C. After incubation for 4 hours, &gt;90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p><b>Functional Testing (15 minute Digest)</b>            A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of pXba DNA and 1 µl of Bsal-HF®v2 incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.</p>	Pass


Assay Name/Specification	Lot # 10177238
<p><b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of pXba DNA and a minimum of 60 units of Bsal-HF@v2 incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	<b>Pass</b>
<p><b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 units of Bsal-HF@v2 incubated for 4 hours at 37°C results in &lt;20% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	<b>Pass</b>
<p><b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] E. coli DNA and a minimum of 100 units of Bsal-HF@v2 incubated for 4 hours at 37°C releases &lt;0.1% of the total radioactivity.</p>	<b>Pass</b>
<p><b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of pXba DNA with Bsal-HF@v2, &gt;95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, &gt;95% can be recut with Bsal-HF@v2.</p>	<b>Pass</b>
<p><b>Protein Purity Assay (SDS-PAGE)</b> Bsal-HF@v2 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Jianying Luo  
Production Scientist  
14 Jul 2022



Michael Tonello  
Packaging Quality Control Inspector  
02 Feb 2023