

## New England Biolabs Certificate of Analysis

**Product Name:** BstYI  
**Catalog Number:** R0523S  
**Concentration:** 10,000 U/ml  
**Unit Definition:** One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in rCutSmart Buffer in 1 hour at 60°C in a total reaction volume of 50 µl.  
**Packaging Lot Number:** 10162339  
**Expiration Date:** 08/2024  
**Storage Temperature:** -20°C  
**Storage Conditions:** 10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @ 25°C)  
**Specification Version:** PS-R0523S v2.0


BstYI Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
R0523SVIAL	BstYI	10162338	Pass
B6004SVIAL	rCutSmart™ Buffer	10161526	Pass

Assay Name/Specification	Lot # 10162339
<b>Non-Specific DNase Activity (16 Hour)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and a minimum of 30 units of BstYI incubated for 16 hours at 60°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>Exonuclease Activity (Radioactivity Release)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [ <sup>3</sup> H] E. coli DNA and a minimum of 50 units of BstYI incubated for 4 hours at 60°C releases <0.1% of the total radioactivity.	Pass
<b>qPCR DNA Contamination (E. coli Genomic)</b> A minimum of 10 units of BstYI is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass
<b>Functional Testing (15 minute Digest)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of Lambda DNA and 1 µl of	Pass

Assay Name/Specification	Lot # 10162339
<p>BstYI incubated for 15 minutes at 60°C results in complete digestion as determined by agarose gel electrophoresis.</p>	
<p><b>Protein Purity Assay (SDS-PAGE)</b> BstYI is &gt;95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.</p>	<b>Pass</b>
<p><b>Ligation and Recutting (Terminal Integrity)</b> After a 20-fold over-digestion of Lambda DNA with BstYI, &gt;95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, &gt;95% can be recut with BstYI.</p>	<b>Pass</b>
<p><b>Endonuclease Activity (Nicking)</b> A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 10 units of BstYI incubated for 4 hours at 60°C results in &lt;50% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.




---

YunJie Sun  
Production Scientist  
17 Aug 2022




---

Erin Varney  
Packaging Quality Control Inspector  
21 Sep 2022