

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Agel
Catalog Number: R0552S
Concentration: 5,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of Lambda DNA in 1 hour at 37°C in a total reaction of 50 μl.

Packaging Lot Number: 10091152
Expiration Date: 03/2022
Storage Temperature: -20°C

Storage Conditions: 250 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50%

Glycerol, 0.15% Triton X-100, 200 µg/ml BSA

Specification Version: PS-R0552S/L v1.0

Agel Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
R0552SVIAL	Agel	10070388	Pass	
B7201SVIAL	NEBuffer™ 1.1	10090429	Pass	
B7024SVIAL	Gel Loading Dye, Purple (6X)	10084974	Pass	

Assay Name/Specification	Lot # 10091152
Blue-White Screening (Terminal Integrity) A sample of LITMUS28i vector linearized with a 10-fold excess of Agel, religated and transformed into an E. coli strain expressing the LacZ beta fragment gene results in <1% white colonies.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 1.1 containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of Agel incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Ligation and Recutting (Terminal Integrity) After a 10-fold over-digestion of Lambda DNA with Agel, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with Agel.	Pass
Non-Specific DNase Activity (16 hour) A 50 µl reaction in NEBuffer 1.1 containing 1 µg of Lambda DNA and a minimum of 5 Units of Agel incubated for 16 hours at 37°C results in a DNA pattern free of	Pass



R0552S / Lot: 10091152

Page 1 of 2

Assay Name/Specification	Lot # 10091152
detectable nuclease degradation as determined by agarose gel electrophoresis. NOTE:	
although no nuclease degradation is detected under these conditions, extended	
incubations and/or high concentrations of this enzyme may result in star activity.	
See the product FAQ for recommended reaction conditions for this enzyme.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang Production Scientist

24 Nov 2020

Josh Hersey

Packaging Quality Control Inspector

24 Nov 2020

