

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: BsrFl-v2
Catalog Number: R0682S
Concentration: 10,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg

of pBR322 DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

Packaging Lot Number: 10092182
Expiration Date: 12/2021
Storage Temperature: -20°C

Storage Conditions: 250 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol,

0.15 % Triton® X-100, 200 μg/ml BSA, (pH 7.4 @ 25°C)

Specification Version: PS-R0682S/L v2.0

BsrFI-v2 Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
R0682SVIAL	BsrFI-v2	10092183	Pass	
B7204SVIAL	CutSmart® Buffer	10091036	Pass	

Assay Name/Specification	Lot # 10092182
Protein Purity Assay (SDS-PAGE) BsrFI-v2 is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Functional Testing (15 minute Digest) A 50 μl reaction in CutSmart® Buffer containing 1 μg of pBR322 DNA and 1 μl of BsrFl-v2 incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in CutSmart® Buffer containing 1 μg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 30 units of BsrFI-v2 incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
<b>Ligation and Recutting (Terminal Integrity)</b> After a 10-fold over-digestion of pBR322 DNA with BsrFI-v2, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BsrFI-v2.	Pass



R0682S / Lot: 10092182

Page 1 of 2



Assay Name/Specification	Lot # 10092182
Non-Specific DNase Activity (16 Hour)	Pass
A 50 µl reaction in CutSmart® Buffer containing 1 µg of pBR322 DNA and a minimum of	
10 units of BsrFI-v2 incubated for 16 hours at 37°C results in a DNA pattern free of	
detectable nuclease degradation as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Penghua Zhang **Production Scientist** 

29 Dec 2020

Michael Tonello

Packaging Quality Control Inspector

29 Dec 2020

