

New England Biolabs Certificate of Analysis

Product Name: Control LAMP Primer Mix (rActin)
Catalog Number: S0164S
Concentration: 10 X Concentrate
Packaging Lot Number: 10159623
Expiration Date: 09/2023
Storage Temperature: -20°C
Specification Version: PS-S0164S v1.0
Composition (1X): Proprietary

Control LAMP Primer Mix (rActin) Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
S0164AVIAL	Internal Control LAMP Primer Mix (rActin)	10154735	Pass

Assay Name/Specification	Lot # 10159623
<p>Endonuclease Activity (Nicking) A 50 µl reaction in NEBuffer 2 containing 1 µg of supercoiled PhiX174 DNA and a minimum of 5 µl of Control LAMP Primer Mix (rActin) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>Functional Testing (LAMP primers) A 25 µl reaction in 1X WarmStart® LAMP Master Mix with UDG in the presence of LAMP Fluorescent Dye and 1X Control LAMP Primer Mix (rActin) containing 10 ng human RNA results in a threshold time of ≤ 20 minutes as determined by fluorescent detection. Reactions that lack human RNA template remain negative over a 30 minute incubation at 65°C.</p>	Pass
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 5 µl of Control LAMP Primer Mix (rActin) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>Phosphatase Activity (pNPP) A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl₂ containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 20 µl of Control LAMP Primer Mix (rActin) incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase</p>	Pass


Assay Name/Specification	Lot # 10159623
<p>activity as determined by spectrophotometric analysis.</p> <p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Control LAMP Primer Mix (rActin) is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	<p>Pass</p>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Christie Vazquez
Production Scientist
12 Aug 2022



Josh Hersey
Packaging Quality Control Inspector
12 Aug 2022