

*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

| Product Name:          | Chitin Resin     |
|------------------------|------------------|
| Catalog Number:        | S6651L           |
| Lot Number:            | 10014180         |
| Expiration Date:       | 06/2021          |
| Storage Temperature:   | 4°C              |
| Specification Version: | PS-S6651S/L v1.0 |

| Chitin Resin Component List |                       |            |                      |
|-----------------------------|-----------------------|------------|----------------------|
| NEB Part Number             | Component Description | Lot Number | Individual QC Result |
| S6651LVIAL                  | Chitin Resin          | 10011011   | Pass                 |

| Assay Name/Specification   | Lot # 10014180 |
|--|----------------|
| <b>Functional Binding Assay (Resin Binding Capacity)</b><br>Chitin Resin (1 ml) was packed into a column and equilibrated with column buffer.<br>Crude extract from E. coli containing a plasmid that expresses a SNAP-intein-chitin<br>binding domain fusion protein (8 ml) was then passed through the column at 25°C,<br>then washed with column buffer and the target protein eluted through cleavage of the<br>fusion protein during overnight incubation with column buffer containing 50 mM DTT<br>at 4°C. Binding capacity was determined to be >2 mg SNAP protein /ml of resin based<br>on A280 of the eluate.  | Pass           |
| <b>Functional Binding Assay (Resin Binding Specificity)</b><br>Chitin Resin (1 ml) was packed into a column and equilibrated with column buffer.<br>Crude extract from E. coli containing a plasmid that expresses a SNAP-intein-chitin<br>binding domain fusion protein (8 ml) was then passed through the column at 25°C,<br>and then washed with column buffer. The target protein was eluted through cleavage<br>of the fusion protein during overnight incubation with column buffer containing 50<br>mM DTT at 4°C. SDS-PAGE of the eluate on a 10-20% Tris-Glycine gel confirms low<br>non-specific binding of extract proteins and high isolation of target. | Pass           |

This product has been tested and shown to be in compliance with all specifications.





*be* INSPIRED *drive* DISCOVERY *stay* GENUINE

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

Brd

Brad Landgraf Production Scientist 21 Jun 2018

Michae -110

Michael Tonello Packaging Quality Control Inspector 26 Jun 2018

