

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Chitin Resin
Catalog Number: S6651S
Lot Number: 10048247
Expiration Date: 03/2022
Storage Temperature: 4°C

Specification Version: PS-S6651S/L v1.0

| Chitin Resin Component List |                       |            |                      |
|-----------------------------|-----------------------|------------|----------------------|
| <b>NEB Part Number</b>      | Component Description | Lot Number | Individual QC Result |
| S6651SVIAL                  | Chitin Resin          | 10037470   | Pass                 |

| Assay Name/Specification   | Lot # 10048247 |  |
|--|----------------|--|
| Functional Binding Assay (Resin Binding Capacity)                                    | Pass           |  |
| Chitin Resin (1 ml) was packed into a column and equilibrated with column buffer.    |                |  |
| Crude extract from E. coli containing a plasmid that expresses a SNAP-intein-chitin  |                |  |
| binding domain fusion protein ( 8 ml ) was then passed through the column at 25°C,   |                |  |
| then washed with column buffer and the target protein eluted through cleavage of the |                |  |
| fusion protein during overnight incubation with column buffer containing 50 mM DTT   |                |  |
| at 4°C. Binding capacity was determined to be >2 mg SNAP protein /ml of resin based  |                |  |
| on A280 of the eluate.   |                |  |
| Functional Binding Assay (Resin Binding Specificity)                                 | Pass           |  |
| Chitin Resin (1 ml) was packed into a column and equilibrated with column buffer.    |                |  |
| Crude extract from E. coli containing a plasmid that expresses a SNAP-intein-chitin  |                |  |
| pinding domain fusion protein (8 ml) was then passed through the column at 25°C,     |                |  |
| and then washed with column buffer. The target protein was eluted through cleavage   |                |  |
| of the fusion protein during overnight incubation with column buffer containing 50   |                |  |
| mM DTT at 4°C. SDS-PAGE of the eluate on a 10-20% Tris-Glycine gel confirms low      |                |  |
| non-specific binding of extract proteins and high isolation of target.               |                |  |

This product has been tested and shown to be in compliance with all specifications.



S6651S / Lot: 10048247

Page 1 of 2

grel 2008

Brad Landgraf Production Scientist 21 Mar 2019 Michael Tonello

Packaging Quality Control Inspector 03 Jul 2019