

New England Biolabs Product Specification

<i>Product Name:</i>	NEBNext [®] End Repair Reaction Buffer
<i>Catalog #:</i>	B6052S
<i>Concentration:</i>	10X Concentrate
<i>Shelf Life:</i>	24 months
<i>Storage Temp:</i>	-20°C
<i>Composition (1X):</i>	50 mM Tris-HCl, 10 mM MgCl ₂ , 10 mM DTT, 1 mM ATP, 0.4 mM dATP, 0.4 mM dCTP, 0.4 mM dGTP, 0.4 mM dTTP, (pH 7.5 @ 25°C)
<i>Specification Version:</i>	PS-B6052S v2.0
<i>Effective Date:</i>	12 Feb 2020

Assay Name/Specification (minimum release criteria)

Endonuclease Activity (Nicking, Buffer) - A 50 µl reaction in 1X NEBNext[®] End Repair Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Non-Specific DNase Activity (16 hour, Buffer) - A 50 µl reaction in 1X NEBNext[®] End Repair Reaction Buffer containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Phosphatase Activity (pNPP, Buffer) - A 200 µl reaction in 1M Diethanolamine @ pH 9.8 and 0.5 mM MgCl₂ containing 2.5 mM *p*-Nitrophenyl Phosphate (pNPP) and a minimum of 20 µl NEBNext[®] End Repair Reaction Buffer incubated for 4 hours at 37°C yields <0.00001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.

RNase Activity (Buffer) - A 10 µl reaction in 1X NEBNext[®] End Repair Reaction Buffer containing 40 ng of a 300 base single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.

One or more products referenced in this document may be covered by a 3rd-party trademark.
Please visit www.neb.com/trademarks for additional information.



Date 12 Feb 2020

Derek Robinson
Director, Quality Control

