

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Product Specification

Product Name:	DNA Polymerase I (E. coli)
Catalog #:	M0209S/L
Concentration:	10,000 units/ml
Unit Definition:	One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid insoluble material in 30 minutes at 37° C.
Shelf Life:	24 months
Storage Temp:	-20°C
Storage Conditions:	25 mM Tris-HCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol, (pH 7.4 @ 25°C)
Specification Version:	PS-M0209S/L v1.0
Effective Date:	04 Aug 2015

Assay Name/Specification (minimum release criteria)

Endonuclease Activity (Nicking) - A 50 μ l reaction in NEBuffer 2 containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 100 units of DNA Polymerase I (*E. coli*) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Phosphatase Activity (pNPP) - A 200 μ l reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl₂ containing 2.5 mM *p*-Nitrophenol Phosphate (pNPP) and a minimum of 100 units DNA Polymerase I (*E. coli*) incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.

Protein Purity Assay (SDS-PAGE) - DNA Polymerase I (*E. coli*) is \geq 99% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.

qPCR DNA Contamination (*E. coli* **Genomic)** - A minimum of 10 units of DNA Polymerase I (*E. coli*) is screened for the presence of *E. coli* genomic DNA using SYBR® Green qPCR with primers specific for the *E. coli* 16S rRNA locus. Results are quantified using a standard curve generated from purified *E. coli* genomic DNA. The measured level of *E. coli* genomic DNA contamination is ≤ 1 *E. coli* genome.

RNase Activity (Extended Digestion) - A 10 μ l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μ l of DNA Polymerase I (*E. coli*) is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.

Derek Robinson Director of Quality Control



Date 04 Aug 2015