

## New England Biolabs Product Specification

<i>Product Name:</i>	<i>Endonuclease VIII</i>
<i>Catalog #:</i>	<i>M0299S/L</i>
<i>Concentration:</i>	<i>10,000 units/ml</i>
<i>Unit Definition:</i>	<i>One unit is defined as the amount of enzyme required to cleave 1 pmol of a 34-mer oligonucleotide duplex containing a single AP site in a total reaction volume of 10 µl in 1 hour at 37°C in 1X Endonuclease VIII Reaction Buffer containing 10 pmol of fluorescently labeled oligonucleotide duplex.</i>
<i>Shelf Life:</i>	<i>12 months</i>
<i>Storage Temp:</i>	<i>-20°C</i>
<i>Storage Conditions:</i>	<i>10 mM Tris-HCl, 250 mM NaCl, 0.1 mM EDTA, 50 % Glycerol, (pH 8.0 @ 25°C)</i>
<i>Specification Version:</i>	<i>PS-M0299S/L v1.0</i>
<i>Effective Date:</i>	<i>19 Mar 2018</i>

### Assay Name/Specification (minimum release criteria)

**Exonuclease Activity (Radioactivity Release)** - A 50 µl reaction in Endonuclease VIII Reaction Buffer containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] *E. coli* DNA and a minimum of 10 units of Endonuclease VIII incubated for 4 hours at 37°C releases <0.5% of the total radioactivity.

**Non-Specific DNase Activity (16 Hour)** - A 50 µl reaction in Endonuclease VIII Reaction Buffer containing 1 µg of Lambda-HindIII DNA and a minimum of 30 units of Endonuclease VIII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

**Protein Purity Assay (SDS-PAGE)** - Endonuclease VIII is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.



Date 19 Mar 2018

Derek Robinson  
Director of Quality Control

