

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Product Specification

Product Name:	Thermostable 5' App DNA/RNA Ligase
Catalog #:	M0319S/L
Concentration:	20 µM
Shelf Life:	24 months
Storage Temp:	-20°C
Storage Conditions:	10 mM Tris-HCl , 50 mM KCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol, (pH 7.4 @ 25°C)
Specification Version:	PS-M0319S/L v1.0
Effective Date:	01 Nov 2018

Assay Name/Specification (minimum release criteria)

Endonuclease Activity (Nicking) - A 50  $\mu$ l reaction in NEBuffer 1 containing 1  $\mu$ g of supercoiled PhiX174 DNA and a minimum of 100 pmol of Thermostable 5' App DNA/RNA Ligase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

**Exonuclease Activity (Radioactivity Release)** - A 50  $\mu$ l reaction in NEBuffer 1 containing 1  $\mu$ g of a mixture of single and doublestranded [<sup>3</sup>H] *E. coli* DNA and a minimum of 100 pmol of Thermostable 5' App DNA/RNA Ligase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

Functional Testing (Targeted Ligation) - A 20  $\mu$ l reaction in 1X NEBuffer 1 containing 20 pmol of 30 bp FAM-labeled single-stranded RNA, 200 pmol 17 bp 5' pre-adenylated single-stranded DNA linker, and 40 pmol Thermostable 5' App DNA/RNA Ligase incubated for 1 hour at 65°C results in ≥80% ligation of the substrate RNA as determined by capillary electrophoresis.

**Phosphatase Activity (pNPP)** - A 200  $\mu$ l reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl<sub>2</sub> containing 2.5 mM *p*-Nitrophenyl Phosphate (pNPP) and a minimum of 100 pmol of Thermostable 5' App DNA/RNA Ligase incubated for 16 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.

**RNase Activity (Extended Digestion)** - A 10  $\mu$ l reaction in NEBuffer 1 containing 40 ng of a 300 base single-stranded RNA and a minimum of 100 pmol of Thermostable 5' App DNA/RNA Ligase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.

Date 01 Nov 2018

Derek Robinson Director of Quality Control



PS-M0319S/L v1.0 Page 1 of 1