240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Product Specification

Product Name: Msz Exonuclease I

Catalog #: M0527S

Concentration: 10,000 units/ml

Unit Definition:

One unit is defined as the amount of enzyme that will catalyze the release of 10 nmol of acid-soluble nucleotide in a total

reaction volume of 100 µl in 15 minutes at 55°C in 1X rCutSmart Buffer with 0.17 mg/ml single-stranded [3H]-DNA

Shelf Life: 24 months
Storage Temp: -20°C

Storage Conditions: 10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, (pH 7.4 @ 25°C)

Specification Version: PS-M0527S v2.0
Effective Date: 10 Aug 2022

Assay Name/Specification (minimum release criteria)

Endonuclease Activity (Circular Single Stranded DNA) - A 50 μ l reaction in rCutSmartTM Buffer containing 1 μ g of M13mp18 Single-stranded DNA and a minimum of 100 units of Msz Exonuclease I incubated for 16 hours at 37°C results in <10% conversion to linear DNA as determined by agarose gel electrophoresis.

Endonuclease Activity (Nicking) - A 50 μ l reaction in rCutSmartTM Buffer containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 100 units of Msz Exonuclease I incubated for 16 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Protein Purity Assay (SDS-PAGE) - Msz Exonuclease I is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.

qPCR DNA Contamination (*E. coli* Genomic) - A minimum of 10 units of Msz Exonuclease I is screened for the presence of *E. coli* genomic DNA using SYBR® Green qPCR with primers specific for the *E. coli* 16S rRNA locus. Results are quantified using a standard curve generated from purified *E. coli* genomic DNA. The measured level of *E. coli* genomic DNA contamination is ≤ 1 *E. coli* genome.

RNase Activity (Extended Digestion) - A 10 μ l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μ l of Msz Exonuclease I is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Derek Robinson Quality Approver

ISO 9001 Registered Quality Management



