

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Product Specification

Product Name:	EcoGII Methyltransferase
Catalog #:	M0603S/L
Concentration:	5,000 units/ml
Unit Definition:	One unit is defined as the amount of enzyme required to completely protect 100 ng of a FAM-labeled dsDNA substrate in 30 minutes at 37°C in a total reaction volume of 20 $\mu$ L against cleavage by MboI restriction endonuclease as determined by capillary electrophoresis.
Shelf Life:	24 months
Storage Temp:	-20°C
Storage Conditions:	250 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 0.15 % Triton X-100, 180 μg/ml BSA, (pH 7.4 @ 25°C)
Specification Version:	<i>PS-M0603S/L v3.0</i>
Effective Date:	09 Sep 2024

Assay Name/Specification (minimum release criteria)

Endonuclease Activity (Nicking) - A 50  $\mu$ l reaction in rCutSmart<sup>TM</sup> Buffer containing 1  $\mu$ g of supercoiled pBR322 DNA and a minimum of 25 units of EcoGII Methyltransferase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

**Exonuclease Activity (Radioactivity Release)** - A 50  $\mu$ l reaction in rCutSmart<sup>TM</sup> Buffer containing 1  $\mu$ g of a mixture of single and double-stranded [<sup>3</sup>H] *E. coli* DNA and a minimum of 25 units of EcoGII Methyltransferase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

Non-Specific DNase Activity (16 Hour) - A 50  $\mu$ l reaction in rCutSmart<sup>TM</sup> Buffer containing 1  $\mu$ g of Lambda DNA and a minimum of 25 units of EcoGII Methyltransferase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

**Protein Purity Assay (SDS-PAGE)** - EcoGII Methyltransferase is  $\geq$  95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.

**RNase Activity (Extended Digestion)** - A 10  $\mu$ l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1  $\mu$ l of EcoGII Methyltransferase is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.



PS-M0603S/L v3.0 Page 1 of 2



240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Product Specification

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit <u>www.neb.com/trademarks</u> for additional information.

Jaurenbran

Lauren Brown Quality Approver



Date 09 Sep 2024