

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Product Specification

Product Name:	Deoxynucleotide (dNTP) Solution Mix
Catalog #:	N0447S/L
Concentration:	10 mM each dNTP
Unit Definition:	N/A
Shelf Life:	24 months
Storage Temp:	-20°C
Storage Conditions:	Supplied in Ultrapure water as a sodium salt (pH 7.5)
Specification Version:	PS-N0447S/L v2.0
Effective Date:	13 Aug 2015

Assay Name/Specification (minimum release criteria)

Endonuclease Activity (Nicking) - A 50 μ l reaction in NEBuffer 2 containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 10 μ l of Deoxynucleotide (dNTP) Solution Mix incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Non-Specific DNase Activity (16 Hour) - A 50 μ l reaction in NEBuffer 2 containing 1 μ g of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 10 μ l of Deoxynucleotide (dNTP) Solution Mix incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

PCR Amplification (0.5 kb Lambda, dNTPs) - A 50 μ l reaction in ThermoPol® Reaction Buffer in the presence of 200 μ M Deoxynucleotide (dNTP) Solution Mix and 0.5 μ M primers containing 1 ng Lambda DNA with 1.25 units of *Taq* DNA Polymerase for 25 cycles of PCR amplification results in the expected 0.5 kb product.

PCR Amplification (2.0 kb Lambda, dNTPs) - A 50 μ l reaction in ThermoPol® Reaction Buffer in the presence of 200 μ M Deoxynucleotide (dNTP) Solution Mix and 0.5 μ M primers containing 1 ng Lambda DNA with 1.25 units of *Taq* DNA Polymerase for 25 cycles of PCR amplification results in the expected 2.0 kb product.

PCR Amplification (5.0 kb Lambda, dNTPs) - A 50 μ l reaction in ThermoPol® Reaction Buffer in the presence of 200 μ M Deoxynucleotide (dNTP) Solution Mix and 0.5 μ M primers containing 1 ng Lambda DNA with 1.25 units of *Taq* DNA Polymerase for 25 cycles of PCR amplification results in the expected 5.0 kb product.

Phosphatase Activity (pNPP) - A 200 μ l reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl₂ containing 2.5 mM *p*-Nitrophenyl Phosphate (pNPP) and a minimum of 40 μ l Deoxynucleotide (dNTP) Solution Mix incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.



PS-N0447S/L v2.0 Page 1 of 2



240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Product Specification

Assay Name/Specification (minimum release criteria)

Physical Purity (HPLC) - Deoxynucleotide (dNTP) Solution Mix is \geq 99% pure as determined by HPLC analysis.

RNase Activity (Extended Digestion) - A 10 μ l reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 μ l of Deoxynucleotide (dNTP) Solution Mix is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.

Date 13 Aug 2015

Derek Robinson Director of Quality Control



PS-N0447S/L v2.0 Page 2 of 2