240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Product Specification

Product Name: DraI

Catalog #: R0129S/L
Concentration: 20,000 units/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in rCutSmart Buffer in 1 hour at 37°C

in a total reaction volume of 50 µl.

Shelf Life: 24 months
Storage Temp: -20°C

Storage Conditions: 10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @ 25°C)

Specification Version: PS-R0129S/L v2.0
Effective Date: 07 Apr 2023

Assay Name/Specification (minimum release criteria)

Exonuclease Activity (Radioactivity Release) - A 50 μ l reaction in rCutSmartTM Buffer containing 1 μ g of a mixture of single and double-stranded [3 H] *E. coli* DNA and a minimum of 200 units of DraI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

Functional Testing (15 minute Digest) - A 50 μl reaction in rCutSmartTM Buffer containing 1 μg of Lambda DNA and 1 μl of DraI incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.

Ligation and Recutting (Terminal Integrity) - After a 20-fold over-digestion of Lambda DNA with DraI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with DraI.

Non-Specific DNase Activity (16 Hour) - A 50 μ l reaction in rCutSmartTM Buffer containing 1 μ g of Lambda DNA and a minimum of 100 units of DraI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Protein Purity Assay (SDS-PAGE) - DraI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.

qPCR DNA Contamination (*E. coli* Genomic) - A minimum of 20 units of DraI is screened for the presence of *E. coli* genomic DNA using SYBR® Green qPCR with primers specific for the *E. coli* 16S rRNA locus. Results are quantified using a standard curve generated from purified *E. coli* genomic DNA. The measured level of *E. coli* genomic DNA contamination is ≤ 1 *E. coli* genome.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Many Gunden

Date 07 Apr 2023

Nancy Considine Quality Approver





