240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350

24 Apr 2013

Date

www.neb.com info@neb.com

New England Biolabs Product Specification

Product Name: XbaI

Catalog #: R0145T/M
Concentration: 100,000 units/ml

Unit Definition:

One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA (dam-/HindIII digest) in 1 hour at 37°C

in a total reaction volume of 50 µl.

Shelf Life: 24 months Storage Temp: $-20 \, ^{\circ}$ C

Storage Conditions: 50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA

Specification Version: PS-R0145T/M v1.0

Effective Date: 24 Apr 2013

Assay Name/Specification (minimum release criteria)

Blue-White Screening (Terminal Integrity) - A sample of pUC19 vector linearized with a 10-fold excess of XbaI, religated and transformed into an *E. coli* strain expressing the LacZ beta fragment gene results in <1% white colonies.

Endonuclease Activity (Nicking) - A 50 μ l reaction in CutSmartTM Buffer containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 100 Units of XbaI incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Exonuclease Activity (Radioactivity Release) - A 50 μ l reaction in CutSmartTM Buffer containing 1 μ g of a mixture of single and double-stranded [3 H] *E. coli* DNA and a minimum of 200 units of XbaI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

Ligation and Recutting (Terminal Integrity) - After a 10-fold over-digestion of Adenovirus-2 DNA with XbaI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with XbaI.

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in CutSmartTM Buffer containing 1 µg of Lambda HindIII dam- DNA and a minimum of 200 Units of XbaI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Protein Purity Assay (SDS-PAGE) - XbaI is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.

Derek Robinson

Director of Quality Control







^{*} The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.