240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Product Specification

Product Name: PvuII

Catalog #: R0151T/M
Concentration: 50,000 units/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in NEBuffer r3.1 in 1 hour at 37°C in

a total reaction volume of 50 µl.

Shelf Life: 24 months
Storage Temp: -20°C

Storage Conditions: 10 mM Tris-HCl, 300 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 500 µg/ml rAlbumin (pH 7.4 @

25°C,

Specification Version: PS-R0151T/M v2.0

Effective Date: 11 Jul 2023

## Assay Name/Specification (minimum release criteria)

Endonuclease Activity (Nicking) - A 50  $\mu$ l reaction in NEBuffer<sup>TM</sup> r3.1 containing 1  $\mu$ g of supercoiled PhiX174 DNA and a minimum of 50 units of PvuII incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Exonuclease Activity (Radioactivity Release) - A 50  $\mu$ l reaction in NEBuffer<sup>TM</sup> r3.1 containing 1  $\mu$ g of a mixture of single and double-stranded [  $^3$ H] *E. coli* DNA and a minimum of 100 units of PvuII incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

Functional Testing (15 minute Digest) - A 50  $\mu$ l reaction in NEBuffer<sup>TM</sup> r3.1 containing 1  $\mu$ g of Lambda DNA and 1  $\mu$ l of PvuII incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.

Ligation and Recutting (Terminal Integrity) - After a 10-fold over-digestion of Lambda DNA with PvuII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with PvuII.

Non-Specific DNase Activity (16 Hour) - A 50  $\mu$ l reaction in NEBuffer<sup>TM</sup> r3.1 containing 1  $\mu$ g of Lambda DNA and a minimum of 10 units of PvuII incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Protein Purity Assay (SDS-PAGE) - PvuII is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit <a href="https://www.neb.com/trademarks">www.neb.com/trademarks</a> for additional information.

Many Gundan

Date 11 Jul 2023

Nancy Considine Quality Approver





