240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Product Specification

Product Name: BstEII
Catalog #: R0162S/L

Concentration: 10,000 units/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 60°C in a total reaction

volume of 50 μ l.

Shelf Life: 24 months Storage Temp: $-20 \,^{\circ}\text{C}$

Storage Conditions: 50 mM KCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA

Specification Version: PS-R0162S/L v1.0

Effective Date: 25 Jun 2013

Assay Name/Specification (minimum release criteria)

Endonuclease Activity (Nicking) - A 50 μ l reaction in NEBuffer 3.1 containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 30 units of BstEII incubated for 4 hours at 60°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Exonuclease Activity (Radioactivity Release) - A 50 μl reaction in NEBuffer 3.1 containing 1 μg of a mixture of single and double-stranded [³H] *E. coli* DNA and a minimum of 50 units of BstEII incubated for 4 hours at 60°C releases <0.1% of the total radioactivity.

Ligation and Recutting (Terminal Integrity) - After a 10-fold over-digestion of Lambda DNA with BstEII, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with BstEII.

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in NEBuffer 3.1 containing 1 µg of Lambda DNA and a minimum of 50 Units of BstEII incubated for 16 hours at 60°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Date

25 Jun 2013

Derek Robinson

Director of Quality Control

nqa。 ISO 9001 Registered Quality





^{*} The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.