240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Product Specification

Product Name: RsaI

Catalog #: R0167S/L
Concentration: 10,000 units/ml

Unit Definition:

One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in rCutSmart™ Buffer in 1 hour at

37°C in a total reaction volume of 50 μl.

Shelf Life: 24 months
Storage Temp: -20°C

Storage Conditions: 10 mM Tris-HCl, 100 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @)

25°C,

Specification Version: PS-R0167S/L v2.0
Effective Date: 03 Mar 2022

## Assay Name/Specification (minimum release criteria)

Exonuclease Activity (Radioactivity Release) - A 50  $\mu$ l reaction in rCutSmart<sup>TM</sup> Buffer containing 1  $\mu$ g of a mixture of single and double-stranded [  $^3$ H] *E. coli* DNA and a minimum of 50 units of RsaI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

Functional Testing (15 minute Digest) - A 50 μl reaction in rCutSmart<sup>TM</sup> Buffer containing 1 μg of Lambda DNA and 1 μl of RsaI incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.

**Ligation and Recutting (Terminal Integrity)** - After a 10-fold over-digestion of Lambda DNA with RsaI, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with RsaI.

Non-Specific DNase Activity (16 Hour) - A 50  $\mu$ l reaction in rCutSmart<sup>TM</sup> Buffer containing 1  $\mu$ g of Lambda DNA and a minimum of 50 units of RsaI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Protein Purity Assay (SDS-PAGE) - RsaI is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.

qPCR DNA Contamination (*E. coli* Genomic) - A minimum of 10 units of RsaI is screened for the presence of *E. coli* genomic DNA using SYBR® Green qPCR with primers specific for the *E. coli* 16S rRNA locus. Results are quantified using a standard curve generated from purified *E. coli* genomic DNA. The measured level of *E. coli* genomic DNA contamination is  $\leq 1$  *E. coli* genome.







240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Product Specification

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit <a href="https://www.neb.com/trademarks">www.neb.com/trademarks</a> for additional information.

Luk Cotumn

Date 03 Mar 2022

Derek Robinson Director, Quality Control





