

## New England Biolabs Product Specification

|                               |   |
|-------------------------------|---|
| <b>Product Name:</b>          | <i>DrdI</i>   |
| <b>Catalog #:</b>             | R0530S/L/V  |
| <b>Concentration:</b>         | 10,000 units/ml   |
| <b>Unit Definition:</b>       | One unit is defined as the amount of enzyme required to digest 1 µg of pUC19 DNA in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 µl. |
| <b>Shelf Life:</b>            | 24 months   |
| <b>Storage Temp:</b>          | -20°C   |
| <b>Storage Conditions:</b>    | 10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @ 25°C)  |
| <b>Specification Version:</b> | PS-R0530S/L/V v4.0  |
| <b>Effective Date:</b>        | 20 Aug 2024   |

### Assay Name/Specification (minimum release criteria)

**Ligation and Recutting (Terminal Integrity)** - After a 10-fold over-digestion of pUC19 DNA with *DrdI*, ~75% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with *DrdI*.

**Exonuclease Activity (Radioactivity Release)** - A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] *E. coli* DNA and a minimum of 50 units of *DrdI* incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

**Functional Testing (15 minute Digest)** - A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of pUC19 DNA and 1 µl of *DrdI* incubated for 15 minutes at 37°C results in complete digestion as determined by agarose gel electrophoresis.

**Non-Specific DNase Activity (16 Hour)** - A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of pUC19 DNA and a minimum of 30 units of *DrdI* incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

**Protein Purity Assay (SDS-PAGE)** - *DrdI* is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.

**qPCR DNA Contamination (*E. coli* Genomic)** - A minimum of 10 units of *DrdI* is screened for the presence of *E. coli* genomic DNA using SYBR® Green qPCR with primers specific for the *E. coli* 16S rRNA locus. Results are quantified using a standard curve generated from purified *E. coli* genomic DNA. The measured level of *E. coli* genomic DNA contamination is ≤ 1 *E. coli* genome.



240 County Road  
Ipswich, MA 01938-2723

Tel 978-927-5054  
Fax 978-921-1350

www.neb.com  
info@neb.com

---

## New England Biolabs Product Specification

*One or more products referenced in this document may be covered by a 3rd-party trademark.  
Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.*

*Nancy Considine*

Date 20 Aug 2024

---

Nancy Considine  
Quality Approver

