

New England Biolabs Product Specification

Product Name:	Nb.BsrDI
Catalog #:	R0648S/L
Concentration:	10,000 units/ml
Unit Definition:	One unit is defined as the amount of enzyme required to convert 1 µg of supercoiled pUC19 DNA to open circular form in rCutSmart Buffer in 1 hour at 37°C in a total reaction volume of 50 µl.
Shelf Life:	24 months
Storage Temp:	-20°C
Storage Conditions:	10 mM Tris-HCl, 200 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml rAlbumin (pH 7.4 @ 25°C)
Specification Version:	PS-R0648S/L v2.0
Effective Date:	20 Aug 2024

Assay Name/Specification (minimum release criteria)

Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of a mixture of single and double-stranded [³H] *E. coli* DNA and a minimum of 100 units of Nb.BsrDI incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in rCutSmart™ Buffer containing 1 µg of pUC19 DNA and a minimum of 30 units of Nb.BsrDI incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

qPCR DNA Contamination (*E. coli* Genomic) - A minimum of 10 units of Nb.BsrDI is screened for the presence of *E. coli* genomic DNA using SYBR® Green qPCR with primers specific for the *E. coli* 16S rRNA locus. Results are quantified using a standard curve generated from purified *E. coli* genomic DNA. The measured level of *E. coli* genomic DNA contamination is ≤ 1 *E. coli* genome.

One or more products referenced in this document may be covered by a 3rd-party trademark.
Please visit www.neb.com/trademarks for additional information.

Nancy Considine

Date 20 Aug 2024

Nancy Considine
Quality Approver

