240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Product Specification

Product Name: EcoRI-HF®

Catalog #: R3101S/L

Concentration: 20,000 units/ml

Unit Definition: One unit is defined as the amount of enzyme required to digest 1 µg of Lambda DNA in 1 hour at 37°C in a total reaction

volume of 50  $\mu$ l.

Shelf Life: 24 months
Storage Temp: -20°C

Storage Conditions: 300 mM NaCl, 10 mM KPO<sub>4</sub>, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 0.15 % TritonX-100, 200 µg/ml BSA,

(pH 7.0 @ 25°C)

Specification Version: PS-R3101S/L v2.0
Effective Date: 16 May 2016

## Assay Name/Specification (minimum release criteria)

Blue-White Screening (Terminal Integrity) - A sample of pUC19 vector linearized with a 10-fold excess of EcoRI-HF<sup>TM</sup>, religated and transformed into an *E. coli* strain expressing the LacZ beta fragment gene results in <1% white colonies.

Endonuclease Activity (Nicking) - A 50  $\mu$ l reaction in CutSmart<sup>TM</sup> Buffer containing 1  $\mu$ g of supercoiled PhiX174 DNA and a minimum of 100 Units of EcoRI-HF<sup>TM</sup> incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Exonuclease Activity (Radioactivity Release) - A 50 μl reaction in CutSmart<sup>TM</sup> Buffer containing 1 μg of a mixture of single and double -stranded [ <sup>3</sup>H] *E. coli* DNA and a minimum of 200 units of EcoRI-HF<sup>TM</sup> incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

**Ligation and Recutting (Terminal Integrity)** - After a 20-fold over-digestion of Lambda DNA with EcoRI-HF<sup>TM</sup>, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with EcoRI-HF<sup>TM</sup>.

Non-Specific DNase Activity (16 Hour) - A 50  $\mu$ l reaction in CutSmart<sup>TM</sup> Buffer containing 1  $\mu$ g of Lambda DNA and a minimum of 100 Units of EcoRI-HF<sup>TM</sup> incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Protein Purity Assay (SDS-PAGE) - EcoRI-HFTM is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.







240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Product Specification

\* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.

Ruh Cotimon

Date 16 May 2016

Derek Robinson Director of Quality Control





