

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Product Specification

Catalog #:R3189MConcentration:100,000 units/mlUnit Definition:One unit is defined as the amount of enzyme required to digest 1 μg of pBC4 DNA in 1 hour at 37°C in a total reaction volume of 50 μl.Shelf Life:24 monthsStorage Temp:-20 °CStorage Conditions:50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSASpecification Version:PS-R3189M v1.0Effective Date:03 May 2013	Product Name:	NotI-HF®
Unit Definition:One unit is defined as the amount of enzyme required to digest 1 μg of pBC4 DNA in 1 hour at 37°C in a total reaction volume of 50 μl.Shelf Life:24 monthsStorage Temp:-20 °CStorage Conditions:50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSASpecification Version:PS-R3189M v1.0	Catalog #:	R3189M
Shelf Life: 24 months Storage Temp: -20 °C Storage Conditions: 50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA Specification Version: PS-R3189M v1.0	Concentration:	100,000 units/ml
Storage Temp: -20 °C Storage Conditions: 50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 µg/ml BSA Specification Version: PS-R3189M v1.0	Unit Definition:	
Storage Conditions:50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSASpecification Version:PS-R3189M v1.0	Shelf Life:	24 months
Specification Version: PS-R3189M v1.0	Storage Temp:	-20 °C
	Storage Conditions:	50 mM NaCl, 10 mM Tris-HCl (pH 7.4), 1 mM DTT, 0.1 mM EDTA, 50% Glycerol, 200 μg/ml BSA
Effective Date: 03 May 2013	Specification Version:	PS-R3189M v1.0
	Effective Date:	03 May 2013

Assay Name/Specification (minimum release criteria)

Endonuclease Activity (Nicking) - A 50 μ l reaction in CutSmartTM Buffer containing 1 μ g of supercoiled PhiX174 DNA and a minimum of 100 Units of NotI-HFTM incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.

Exonuclease Activity (Radioactivity Release) - A 50 μ l reaction in CutSmartTM Buffer containing 1 μ g of a mixture of single and doublestranded [³H] *E. coli* DNA and a minimum of 200 units of NotI-HFTM incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

Ligation and Recutting (Terminal Integrity) - After a 10-fold over-digestion of pBC4 DNA with NotI-HFTM, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with NotI-HFTM.

Non-Specific DNase Activity (16 Hour) - A 50 μ l reaction in CutSmartTM Buffer containing 1 μ g of pBC4 DNA and a minimum of 200 Units of NotI-HFTM incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.

Derek Robinson Director of Quality Control



Date 03 May 2013