240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350

Date

www.neb.com info@neb.com

## New England Biolabs Product Specification

Product Name: EagI-HF®

Catalog #: R3505S/L

Concentration: 20,000 units/ml

Unit Definition:

One unit is defined as the amount of enzyme required to digest 1 µg of pXba DNA in 1 hour at 37°C in a total reaction

volume of 50 µl.

Shelf Life: 24 months
Storage Temp: -20°C

Storage Conditions: 500 mM NaCl, 10 mM Tris-HCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 200 μg/ml BSA, (pH 7.4 @ 25°C)

Specification Version: PS-R3505S/L v2.0
Effective Date: 11 May 2016

## Assay Name/Specification (minimum release criteria)

Blue-White Screening (Terminal Integrity) - A sample of Litmus 38i vector linearized with a 10-fold excess of EagI-HF<sup>TM</sup>, religated and transformed into an  $E.\ coli$  strain expressing the LacZ beta fragment gene results in <1% white colonies.

Endonuclease Activity (Nicking) - A 50  $\mu$ l reaction in CutSmart<sup>TM</sup> Buffer containing 1  $\mu$ g of supercoiled PhiX174 DNA and a minimum of 20 Units of EagI-HF<sup>TM</sup> incubated for 4 hours at 37°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.

Exonuclease Activity (Radioactivity Release) - A 50 μl reaction in CutSmart<sup>TM</sup> Buffer containing 1 μg of a mixture of single and double -stranded [ <sup>3</sup>H] *E. coli* DNA and a minimum of 100 units of EagI-HF<sup>TM</sup> incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.

Ligation and Recutting (Terminal Integrity) - After a 20-fold over-digestion of pXba DNA with EagI-HF<sup>TM</sup>, >95% of the DNA fragments can be ligated with T4 DNA ligase in 16 hours at 16°C. Of these ligated fragments, >95% can be recut with EagI-HF<sup>TM</sup>.

Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in CutSmart<sup>TM</sup> Buffer containing 1 µg of pXba DNA and a minimum of 100 Units of EagI-HF<sup>TM</sup> incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.

Protein Purity Assay (SDS-PAGE) - EagI-HF<sup>TM</sup> is >95% pure as determined by SDS PAGE analysis using Coomassie Blue detection.

Derek Robinson

Director of Quality Control







11 May 2016

<sup>\*</sup> The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.