

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Product Specification

Product Name:	Monarch® Genomic DNA Purification Kit
Catalog #:	T3010S/L
Kit Components:	Monarch® gDNA Tissue Lysis Buffer (T3011) Monarch® gDNA Cell Lysis Buffer (T3012) Monarch® gDNA Blood Lysis Buffer (T3013) Monarch® gDNA Binding Buffer (T3014) Monarch® gDNA Wash Buffer (T3015) Monarch® gDNA Elution Buffer (T3016) Monarch® gDNA Purification Columns (T3017) Proteinase K, Molecular Biology Grade (P8107) Monarch® RNase A (T3018) Monarch® Collection Tubes II (T2018)
Shelf Life:	18 months
Storage Temp:	25°C
Specification Version:	PS-T3010S/L v1.0
Effective Date:	25 Apr 2019

Assay Name/Specification (minimum release criteria)

Functional Testing (Blood Cell Lysis Buffer, Monarch[®]) - Genomic DNA is purified from 8 samples of whole pig blood using the Protocol for Extraction and Purification of Genomic DNA from Mammalian Whole Blood (non-nucleated), with $\geq 2 \mu g$ of gDNA being recovered in $\geq 80\%$ of the samples. OD 260/280 and 260/230 are ≥ 1.75 in $\geq 80\%$ of the samples.

Functional Testing (Cell Lysis Buffer, Monarch[®]) - Genomic DNA is purified from 8 individual aliquots of HeLa cells (10^6 cells/aliquot) using the Protocol for Extraction and Purification of Genomic DNA from Cultured Cells, resulting in $\ge 3 \mu g$ of gDNA being recovered in $\ge 80\%$ of the samples. OD 260/280 and 260/230 are ≥ 1.75 in $\ge 80\%$ of the samples.

Functional Testing (DNA Recovery and Purity) - Twenty-four Monarch® gDNA Purification Columns are tested with 5 μ g of HindIII digested Lambda resulting in \geq 75% recovery in \geq 80% of the samples. OD 260/280 and 260/230 are \geq 1.75 in \geq 80% of the samples.

Functional Testing (RNase A, Monarch[®]) - A 10 μ l reaction in NEBuffer 4 containing 40 ng of fluorescein labeled RNA transcript and RNase A is incubated at 37°C. After incubation for 5 minutes, complete disappearance of the RNA substrate occurs at $\leq 1.0 \ \mu$ g/ml of RNase A, as determined by gel electrophoresis using fluorescent detection.



PS-T3010S/L v1.0 Page 1 of 2



240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350

www.neb.com info@neb.com

New England Biolabs Product Specification

Assay Name/Specification (minimum release criteria)

Functional Testing (Tissue Lysis Buffer, Monarch®) - Genomic DNA is purified from 8 NEB10-Beta samples treated using the Supplemental Protocol for Genomic DNA Purification from Gram-negative Bacteria, resulting in $\geq 5 \ \mu g$ of gDNA being recovered in $\geq 80\%$ of the samples. OD 260/280 and 260/230 are ≥ 1.75 in $\geq 80\%$ of the samples.

* Individual Product Component Note - Standard Quality Control Tests are performed for each component included in Monarch® Genomic DNA Purification Kit and meet the designated specifications.

Date 25 Apr 2019

Derek Robinson Director of Quality Control



PS-T3010S/L v1.0 Page 2 of 2