240 County Road Ipswich, MA 01938-2723

Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Product Specification

Product Name: Monarch® Genomic DNA Purification Kit

*Catalog #: T3010S/L* 

Kit Components: Monarch® gDNA Tissue Lysis Buffer (T3011)

Monarch® gDNA Cell Lysis Buffer (T3012)
Monarch® gDNA Blood Lysis Buffer (T3013)
Monarch® gDNA Binding Buffer (T3014)
Monarch® gDNA Wash Buffer (T3015)
Monarch® gDNA Elution Buffer (T3016)
Monarch® gDNA Purification Columns (T3017)
Proteinase K, Molecular Biology Grade (P8107)

Monarch® RNase A (T3018)

Monarch® Collection Tubes II (T2018)

Shelf Life: 24 months
Storage Temp: 25°C

Specification Version: PS-T3010S/L v2.0
Effective Date: 16 May 2022

## Assay Name/Specification (minimum release criteria)

Functional Testing (Blood Cell Lysis Buffer, Monarch®) - Genomic DNA is purified from 8 samples of whole pig blood using the Protocol for Extraction and Purification of Genomic DNA from Mammalian Whole Blood (non-nucleated), with  $\geq 2 \mu g$  of gDNA being recovered in  $\geq 80\%$  of the samples. OD 260/280 and 260/230 are  $\geq 1.75$  in  $\geq 80\%$  of the samples.

Functional Testing (Cell Lysis Buffer, Monarch®) - Genomic DNA is purified from 8 individual aliquots of HeLa cells (  $10^6$  cells/aliquot) using the Protocol for Extraction and Purification of Genomic DNA from Cultured Cells, resulting in  $\geq 3 \mu g$  of gDNA being recovered in  $\geq 80\%$  of the samples. OD 260/280 and 260/230 are  $\geq 1.75$  in  $\geq 80\%$  of the samples.

Functional Testing (DNA Recovery and Purity) - Twenty-four Monarch® gDNA Purification Columns are tested with 5  $\mu$ g of HindIII digested Lambda resulting in ≥75% recovery in ≥80% of the samples. OD 260/280 and 260/230 are ≥1.75 in ≥80% of the samples.

Functional Testing (RNase A, Monarch®) - A 10  $\mu$ l reaction in NEBuffer 4 containing 40 ng of fluorescein labeled RNA transcript and RNase A is incubated at 37°C. After incubation for 5 minutes, complete disappearance of the RNA substrate occurs at  $\leq$ 1.0  $\mu$ g/ml of RNase A, as determined by gel electrophoresis using fluorescent detection.







240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Product Specification

Assay Name/Specification (minimum release criteria)

Functional Testing (Tissue Lysis Buffer, Monarch®) - Genomic DNA is purified from 8 NEB10-Beta samples treated using the Supplemental Protocol for Genomic DNA Purification from Gram-negative Bacteria, resulting in  $\geq$ 5 µg of gDNA being recovered in  $\geq$ 80% of the samples. OD 260/280 and 260/230 are  $\geq$ 1.75 in  $\geq$ 80% of the samples.

\* Individual Product Component Note - Standard Quality Control Tests are performed for each component included in Monarch® Genomic DNA Purification Kit and meet the designated specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit <a href="https://www.neb.com/trademarks">www.neb.com/trademarks</a> for additional information.

Nuk Cotum

Date 16 May 2022

Director, Quality Control

Derek Robinson





