

## New England Biolabs Certificate of Analysis

**Product Name:** NEBNext<sup>®</sup> dsDNA Fragmentase<sup>®</sup> Reaction Buffer v2  
**Catalog Number:** B0349S  
**Concentration:** 10 X Concentrate  
**Packaging Lot Number:** 10161450  
**Expiration Date:** 01/2024  
**Storage Temperature:** -20°C  
**Specification Version:** PS-B0349S v2.0  
**Composition (1X):** 20 mM Tris-HCl, 15 mM MgCl<sub>2</sub>, 50 mM NaCl, 0.15% Triton<sup>®</sup>X-100, 0.1 mg/ml BSA, (pH 7.5 @ 25°C)

NEBNext <sup>®</sup> dsDNA Fragmentase <sup>®</sup> Reaction Buffer v2 Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
B0349SVIAL	NEBNext <sup>®</sup> dsDNA Fragmentase <sup>®</sup> Reaction Buffer v2	10133265	Pass

Assay Name/Specification	Lot # 10161450
<b>Non-Specific DNase Activity (16 hour, Buffer)</b> A 50 µl reaction in 1X NEBNext <sup>®</sup> dsDNA Fragmentase <sup>®</sup> Reaction Buffer v2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
<b>Phosphatase Activity (pNPP, Buffer)</b> A 200 µl reaction in 1M Diethanolamine @ pH 9.8 and 0.5 mM MgCl <sub>2</sub> containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 20 µl NEBNext <sup>®</sup> dsDNA Fragmentase <sup>®</sup> Reaction Buffer v2 incubated for 4 hours at 37°C yields <0.00001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
<b>Protease Activity (SDS-PAGE, Buffer)</b> A 30 µl reaction in 1X NEBNext <sup>®</sup> dsDNA Fragmentase <sup>®</sup> Reaction Buffer v2 incubated with 24 µg of a standard mixture of proteins for 20 hours at 37°C resulted in no proteolytic activity detected by SDS-PAGE.	Pass
<b>Endonuclease Activity (Nicking, Buffer)</b> A 50 µl reaction in 1X NEBNext <sup>®</sup> dsDNA Fragmentase <sup>®</sup> Reaction Buffer v2 containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



---

Christine Sumner  
Production Scientist  
05 Aug 2022



---

Michael Tonello  
Packaging Quality Control Inspector  
05 Aug 2022