

## New England Biolabs Certificate of Analysis

**Product Name:** *Isothermal Amplification Buffer II Pack*  
**Catalog Number:** *B0374S*  
**Concentration:** *10 X Concentrate*  
**Packaging Lot Number:** *10251143*  
**Expiration Date:** *08/2026*  
**Storage Temperature:** *-20°C*  
**Specification Version:** *PS-B0374S v2.0*  
**Composition (1X):** *20 mM Tris-HCl, 10 mM (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub>, 150 mM KCl, 2 mM MgSO<sub>4</sub>, 0.1 % Tween® 20, (pH 8.8 @ 25°C)*


Isothermal Amplification Buffer II Pack Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
B0374SVIAL	Isothermal Amplification Buffer II Pack	10204644	Pass

Assay Name/Specification	Lot # 10251143
<b>Endonuclease Activity (Nicking, Buffer)</b> A 50 µl reaction in 2X Isothermal Amplification Buffer II containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>Non-Specific DNase Activity (16 hour, Buffer)</b> A 50 µl reaction in 2X Isothermal Amplification Buffer II containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	<b>Pass</b>
<b>Phosphatase Activity (pNPP, Buffer)</b> A 200 µl reaction in 1M Diethanolamine @ pH 9.8 and 0.5 mM MgCl <sub>2</sub> containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 40 µl 10X Isothermal Amplification Buffer II incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	<b>Pass</b>
<b>RNase Activity Assay (4 Hour Digestion)</b> A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Isothermal Amplification Buffer II incubated for 4 hours at 37°C results in no detectable degradation of the RNA as determined by gel electrophoresis using fluorescent detection.	<b>Pass</b>

Assay Name/Specification	Lot # 10251143
<p><b>pH (buffers/solutions)</b> The pH of 10X Isothermal Amplification Buffer II is between pH 8.7 and 8.9 at 25°C.</p>	<b>Pass</b>
<p><b>qPCR DNA Contamination (E. coli Genomic, Buffer)</b> A minimum of 1 µl of Isothermal Amplification Buffer II is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.



Lea Antonopoulos  
Production Scientist  
19 Sep 2023



Michael Tonello  
Packaging Quality Control Inspector  
23 Jul 2024