

New England Biolabs Certificate of Analysis

Product Name: *Isothermal Amplification Buffer*
Catalog Number: *B0537S*
Concentration: *10 X Concentrate*
Packaging Lot Number: *10254032*
Expiration Date: *02/2027*
Storage Temperature: *-20°C*
Specification Version: *PS-B0537S v2.0*
Composition (1X): *20 mM Tris-HCl, 50 mM KCl, 10 mM (NH₄)₂SO₄, 2 mM MgSO₄, 0.1 % Tween® 20, (pH 8.8 @ 25°C)*


Isothermal Amplification Buffer Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
B0537SVIAL	Isothermal Amplification Buffer	10231027	Pass

Assay Name/Specification	Lot # 10254032
<p>Endonuclease Activity (Nicking, Buffer) A 50 µl reaction in 2X Isothermal Amplification Buffer containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 2X Isothermal Amplification Buffer containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>Phosphatase Activity (pNPP, Buffer) A 200 µl reaction in 1M Diethanolamine @ pH 9.8 and 0.5 mM MgCl₂ containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 40 µl Isothermal Amplification Buffer incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p>	Pass
<p>RNase Activity Assay (4 Hour Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Isothermal Amplification Buffer is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass

Assay Name/Specification	Lot # 10254032
<p>pH (buffers/solutions) The pH of 10X Isothermal Amplification Buffer is between pH 8.7 and 8.9 at 25°C.</p>	Pass
<p>qPCR DNA Contamination (E. coli Genomic, Buffer) A minimum of 1 µl of Isothermal Amplification Buffer is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.


 Lea Antonopoulos
 Production Scientist
 01 Apr 2024


 Talia Monkiewicz
 Packaging Quality Control Inspector
 09 Aug 2024