

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: CutSmart® Buffer

Catalog Number: B7204S

Concentration: 10 X Concentrate

Lot Number: 10048294
Expiration Date: 05/2022
Storage Temperature: -20°C

Specification Version: PS-B7204S v1.0

Composition (1X): 50 mM Potassium Acetate , 20 mM Tris Acetate, 10 mM Magnesium

Acetate, 100 μg/ml BSA, (pH 7.9 @ 25°C)

CutSmart® Buffer Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
B7204SVIAL	CutSmart® Buffer	10043913	Pass	

Assay Name/Specification	Lot # 10048294
Conductivity (buffers/solutions)	Pass
The conductivity of 10X CutSmart® Buffer is between 40 and 46 mS at 25°C.	
En la manda de la Activita (Alialian Boffee)	<b>D</b>
Endonuclease Activity (Nicking, Buffer)	Pass
A 50 μl reaction in 1X CutSmart® Buffer containing 1 μg of supercoiled PhiX174 DNA	
incubated for 4 hours at 37°C results in <10% conversion to the nicked form as	
determined by agarose gel electrophoresis.	
Functional Testing (Restriction Digest, Buffer)	Pass
A 50 µl reaction in 1X CutSmart® Buffer containing 1 µg of Lambda dam- DNA and 1	
unit of Clal incubated for 1 hour at 37°C results in complete digestion of the	
substrate DNA as determined by agarose gel electrophoresis.	
Functional Testing (Restriction Digest, Buffer)	Pass
A 50 µl reaction in 1X CutSmart® Buffer containing 1 µg of Lambda DNA and 1 unit of	Fass
Mscl incubated for 1 hour at 37°C results in complete digestion of the substrate DNA	
as determined by agarose gel electrophoresis.	
as determined by againess generosites	
Non-Specific DNase Activity (16 hour, Buffer)	Pass
A 50 μl reaction in 1X CutSmart® Buffer containing 1 μg of PhiX174-HaeIII DNA	
incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease	
degradation as determined by agarose gel electrophoresis.	



B7204S / Lot: 10048294

This product has been tested and shown to be in compliance with all specifications.

single-stranded RNA is incubated at 37°C. After incubation for 16 hours, >90% of the

substrate RNA remains intact as determined by fluorescent detection.

Dreen Duquette Production Scientist

28 May 2019

Mary Conflon

Packaging Quality Control Inspector

19 Jun 2019



B7204S / Lot: 10048294

Page 2 of 2