

New England Biolabs Certificate of Analysis


Product Name: ThermoPol® Reaction Buffer Pack
Catalog Number: B9004S
Concentration: 10 X Concentrate
Lot Number: 10048538
Expiration Date: 08/2023
Storage Temperature: -20°C
Specification Version: PS-B9004S v1.0
Composition (1X): 20 mM Tris-HCl, 10 mM (NH₄)₂SO₄, 10 mM KCl, 2 mM MgSO₄, 0.1 % Triton®X-100, (pH 8.8 @ 25°C)

ThermoPol® Reaction Buffer Pack Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
B9004SVIAL	ThermoPol® Reaction Buffer Pack	10041932	Pass
B1003SVIAL	Magnesium Sulfate (MgSO ₄) Solution	10042724	Pass

Assay Name/Specification	Lot # 10048538
Endonuclease Activity (Nicking, Buffer) A 50 µl reaction in 2X ThermoPol® Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Non-Specific DNase Activity (16 hour, Buffer) A 50 µl reaction in 2X ThermoPol® Reaction Buffer containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	Pass
PCR Amplification (5 kb Lambda DNA, Buffer) A 50 µl reaction in ThermoPol® Reaction Buffer in the presence of 200 µM dNTPs and 0.2 µM primers containing 5 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 5 kb product.	Pass
pH (buffers/solutions) The pH of 10X ThermoPol® Reaction Buffer is between pH 8.7 and 8.9 at 25°C.	Pass
Phosphatase Activity (pNPP, Buffer) A 200 µl reaction in 1M Diethanolamine @ pH 9.8 and 0.5 mM MgCl ₂ containing 2.5 mM	Pass

Assay Name/Specification	Lot # 10048538
<p>p-Nitrophenyl Phosphate (pNPP) and a minimum of 40 µl ThermoPol® Reaction Buffer incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p>	
<p>qPCR DNA Contamination (E. coli Genomic, Buffer) A minimum of 1 µl of ThermoPol® Reaction Buffer is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of ThermoPol® Reaction Buffer is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.



Doreen Duquette
Production Scientist
18 Apr 2019



Jay Minichiello
Packaging Quality Control Inspector
09 Aug 2019