

## New England Biolabs Certificate of Analysis


**Product Name:** T7 Express lysY/lq Competent *E. coli* (High Efficiency)  
**Catalog Number:** C3013I  
**Lot Number:** 10030705  
**Expiration Date:** 09/2020  
**Storage Temperature:** -80°C  
**Specification Version:** PS-C3013I v1.0

T7 Express lysY/lq Competent <i>E. coli</i> (High Efficiency) Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
C3013IVIAL	T7 Express lysY/lq Competent <i>E. coli</i> (High Efficiency)	10017089	Pass


Assay Name/Specification	Lot # 10030705
<b>Antibiotic Resistance (Chloramphenicol)</b> 15 µl of untransformed T7 Express lysY/lq Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Chloramphenicol will form colonies after incubation for 16 hours at 37°C.	Pass
<b>Antibiotic Resistance (Nitrofurantoin)</b> 15 µl of untransformed T7 Express lysY/lq Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Nitrofurantoin will form colonies after incubation for 16 hours at 37°C.	Pass
<b>Antibiotic Sensitivity (Ampicillin)</b> 15 µl of untransformed T7 Express lysY/lq Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Ampicillin will not form colonies after incubation for 16 hours at 37°C.	Pass
<b>Antibiotic Sensitivity (Kanamycin)</b> 15 µl of untransformed T7 Express lysY/lq Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Kanamycin will not form colonies after incubation for 16 hours at 37°C.	Pass
<b>Antibiotic Sensitivity (Spectinomycin)</b> 15 µl of untransformed T7 Express lysY/lq Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Spectinomycin will not form colonies after incubation for 16 hours at 37°C.	Pass

Assay Name/Specification	Lot # 10030705
<p><b>Antibiotic Sensitivity (Streptomycin)</b> 15 µl of untransformed T7 Express lysY/lq Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Streptomycin will not form colonies after incubation for 16 hours at 37°C.</p>	<b>Pass</b>
<p><b>Antibiotic Sensitivity (Tetracycline)</b> 15 µl of untransformed T7 Express lysY/lq Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Tetracycline will not form colonies after incubation for 16 hours at 37°C.</p>	<b>Pass</b>
<p><b>Phage Resistance (φ 80)</b> 15 µl of untransformed T7 Express lysY/lq Competent E. coli (High Efficiency) streaked onto a Rich Broth plate does not support plaque formation by phage φ 80 after incubation for 16 hours at 37°C.</p>	<b>Pass</b>
<p><b>Transformation Efficiency</b> 50 µl of T7 Express lysY/lq Competent E. coli (High Efficiency) cells were transformed with 100 pg of pUC19 DNA using the transformation protocol provided. Incubation overnight on LB-Ampicillin plates at 37°C resulted in &gt;0.6 x 10<sup>9</sup> cfu/µg of DNA.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.



Lixin An  
Production Scientist  
14 Sep 2018



Corey Rabeau  
Packaging Quality Control Inspector  
28 Nov 2018