

New England Biolabs Certificate of Analysis


Product Name: NEB[®] 10-beta Competent *E. coli* (High Efficiency)
 Catalog Number: C3019I
 Lot Number: 10034344
 Expiration Date: 01/2020
 Storage Temperature: -80°C
 Specification Version: PS-C3019H/I v1.0

NEB [®] 10-beta Competent <i>E. coli</i> (High Efficiency) Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
N3041AVIAL	pUC19 Vector	10032414	Pass
C3019IVIAL	NEB [®] 10-beta Competent <i>E. coli</i> (High Efficiency)	10029632	Pass
B9035SVIAL	NEB [®] 10-beta/Stable Outgrowth Medium	10018488	Pass

Assay Name/Specification	Lot # 10034344
Antibiotic Sensitivity (Nitrofurantoin) 15 µl of untransformed NEB [®] 10-beta Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Nitrofurantoin will not form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Sensitivity (Spectinomycin) 15 µl of untransformed NEB [®] 10-beta Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Spectinomycin will not form colonies after incubation for 16 hours at 37°C.	Pass
Antibiotic Sensitivity (Tetracycline) 15 µl of untransformed NEB [®] 10-beta Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Tetracycline will not form colonies after incubation for 16 hours at 37°C.	Pass
Blue-White Screening (α-complementation, Competent Cells) NEB [®] 10-beta Competent <i>E. coli</i> (High Efficiency) were shown to be suitable for blue/white screening by α-complementation of the β-galactosidase gene using pUC19.	Pass
Phage Resistance (φ 80) 15 µl of untransformed NEB [®] 10-beta Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate does not support plaque formation by phage φ 80 after incubation for 16 hours at 37°C.	Pass

Assay Name/Specification	Lot # 10034344
<p>Transformation Efficiency 50 µl of NEB® 10-beta Competent E. coli (High Efficiency) cells were transformed with 100 pg of pUC19 DNA using the transformation protocol provided. Incubation overnight on LB-Ampicillin plates at 37°C resulted in >1 x 10e9 cfu/µg of DNA.</p>	Pass
<p>Antibiotic Resistance (Streptomycin) 15 µl of untransformed NEB® 10-beta Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Streptomycin will form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Sensitivity (Ampicillin) 15 µl of untransformed NEB® 10-beta Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Ampicillin will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Sensitivity (Chloramphenicol) 15 µl of untransformed NEB® 10-beta Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Chloramphenicol will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Sensitivity (Kanamycin) 15 µl of untransformed NEB® 10-beta Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Kanamycin will not form colonies after incubation for 16 hours at 37°C.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.



Lixin An
Production Scientist
21 Dec 2018



Nick Privitera
Packaging Quality Control Inspector
18 Jan 2019