

New England Biolabs Certificate of Analysis

Product Name: NEB[®] 10-beta Competent *E. coli* (High Efficiency)
 Catalog Number: C3019P
 Packaging Lot Number: 10179582
 Expiration Date: 12/2023
 Storage Temperature: -80°C
 Specification Version: PS-C3019P v1.0

NEB [®] 10-beta Competent <i>E. coli</i> (High Efficiency) Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
N3041AVIAL	pUC19 Vector	10173306	Pass
C3019PVIAL	NEB 10-beta Competent <i>E. coli</i> (High Efficiency)	10173371	Pass
B9035SVIAL	NEB [®] 10-beta/Stable Outgrowth Medium	10166802	Pass

Assay Name/Specification	Lot # 10179582
<p>Antibiotic Sensitivity (Ampicillin) 15 µl of untransformed NEB[®] 10-beta Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Ampicillin will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Resistance (Streptomycin) 15 µl of untransformed NEB[®] 10-beta Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Streptomycin will form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Sensitivity (Chloramphenicol) 15 µl of untransformed NEB[®] 10-beta Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate containing Chloramphenicol will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Phage Resistance (φ 80) 15 µl of untransformed NEB[®] 10-beta Competent <i>E. coli</i> (High Efficiency) streaked onto a Rich Broth plate does not support plaque formation by phage φ 80 after incubation for 16 hours at 37°C.</p>	Pass
<p>Blue-White Screening (α-complementation, Competent Cells) NEB[®] 10-beta Competent <i>E. coli</i> (High Efficiency) were shown to be suitable for blue/white screening by α-complementation of the β-galactosidase gene using pUC19.</p>	Pass


Assay Name/Specification	Lot # 10179582
<p>Transformation Efficiency 20 µl of NEB® 10-beta Competent E. coli (High Efficiency) cells were transformed with 100 pg of pUC19 DNA using the transformation protocol provided. Incubation overnight on LB-Ampicillin plates at 37°C resulted in >1 x 10e8 cfu/µg of DNA.</p>	Pass
<p>Antibiotic Sensitivity (Tetracycline) 15 µl of untransformed NEB® 10-beta Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Tetracycline will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Sensitivity (Spectinomycin) 15 µl of untransformed NEB® 10-beta Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Spectinomycin will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Sensitivity (Kanamycin) 15 µl of untransformed NEB® 10-beta Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Kanamycin will not form colonies after incubation for 16 hours at 37°C.</p>	Pass
<p>Antibiotic Sensitivity (Nitrofurantoin) 15 µl of untransformed NEB® 10-beta Competent E. coli (High Efficiency) streaked onto a Rich Broth plate containing Nitrofurantoin will not form colonies after incubation for 16 hours at 37°C.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Lixin An
Production Scientist
02 Dec 2022



Anna Sorensen
Packaging Quality Control Inspector
31 Jan 2023