

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Phusion® High-Fidelity PCR Kit

Catalog Number: E0553S
Lot Number: 10034491
Expiration Date: 10/2020
Storage Temperature: -20°C

Specification Version: PS-E0553S/L v1.0

Phusion® High-Fidelity PCR Kit Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
S0536AVIAL	10 kb Control Primer Mix	10023562	Pass
S0535AVIAL	1.3 kb Control Primer Mix	10023561	Pass
N3010AVIAL	Control Lambda Template	10022938	Pass
N0447AVIAL	Deoxynucleotide (dNTP) Solution Mix	10022939	Pass
N0303AVIAL	Quick-Load® DNA Marker, Broad Range	10022786	Pass
M0530AVIAL	Phusion® High-Fidelity DNA Polymerase	10021635	Pass
B0519SVIAL	Phusion® GC Buffer Pack	0051804	Pass
B0518SVIAL	Phusion® HF Buffer Pack	0071804	Pass
B0515AVIAL	DMSO	10019997	Pass
B0510AVIAL	MgCl2 Solution (50 mM)	10033766	Pass

Assay Name/Specification	Lot # 10034491
* Individual Product Component Note	Pass
Standard Quality Control Tests are performed for each component included in Phusion® High-Fidelity PCR Kit and meet the designated specifications.	
PCR Amplification (7.5 kb Human Genomic DNA) A 50 μl reaction in Phusion® HF Buffer in the presence of 200 μM dNTPs and 1 μM primers containing 50 ng Human Genomic DNA with 1 unit of Phusion® DNA Polymerase for 30 cycles of PCR amplification results in the expected 7.5 kb product.	Pass
PCR Amplification (20 kb Lambda DNA) A 50 μl reaction in Phusion® HF Buffer in the presence of 200 μM dNTPs and 1 μM primers containing 10 ng Lambda DNA with 1 unit of Phusion® DNA Polymerase for 22 cycles of PCR amplification results in the expected 20 kb product.	Pass
Endonuclease Activity (Nicking, Polymerase, dNTP)	Pass



E0553S / Lot: 10034491

Page 1 of 2



Assay Name/Specification	Lot # 10034491
A 50 μl reaction in NEBuffer 2 in the presence of 200 μM dNTPs containing 1 μg of	
supercoiled PhiX174 DNA and a minimum of 10 units of Phusion® DNA Polymerase	
incubated for 4 hours at either 37°C or 72°C results in <10% conversion to the	
nicked form as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

Lynne Apone **Production Scientist** 10 Oct 2018

Josh Hersey Packaging Quality Control Inspector

22 Jan 2019

