

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Phusion® High-Fidelity PCR Kit

Catalog Number: E0553S
Packaging Lot Number: 10086370
Expiration Date: 08/2022
Storage Temperature: -20°C

Specification Version: PS-E0553S/L v1.0

Phusion® High-Fidelity PCR Kit Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
S0536AVIAL	10 kb Control Primer Mix	10082381	Pass
S0535AVIAL	1.3 kb Control Primer Mix	10082382	Pass
N3010AVIAL	Control Lambda Template	10082380	Pass
N0447AVIAL	Deoxynucleotide (dNTP) Solution Mix	10084014	Pass
N0303AVIAL	Quick-Load® DNA Marker, Broad Range	10086618	Pass
M0530AVIAL	Phusion® High-Fidelity DNA Polymerase	10084004	Pass
B0519SVIAL	Phusion® GC Buffer Pack	10073670	Pass
B0518SVIAL	Phusion® HF Buffer Pack	10073662	Pass
B0515AVIAL	DMSO	10081387	Pass
B0510AVIAL	MgCl2 Solution (50 mM)	10073655	Pass

Assay Name/Specification	Lot # 10086370
* Individual Product Component Note Standard Quality Control Tests are performed for each component included in Phusion®	Pass
High-Fidelity PCR Kit and meet the designated specifications.	
PCR Amplification (7.5 kb Human Genomic DNA)	Pass
A 50 μl reaction in Phusion® HF Buffer in the presence of 200 μM dNTPs and 1 μM primers containing 50 ng Human Genomic DNA with 1 unit of Phusion® DNA Polymerase for 30 cycles of PCR amplification results in the expected 7.5 kb product.	
PCR Amplification (20 kb Lambda DNA)	Pass
A 50 µl reaction in Phusion® HF Buffer in the presence of 200 µM dNTPs and 1 µM primers containing 10 ng Lambda DNA with 1 unit of Phusion® DNA Polymerase for 22 cycles of PCR amplification results in the expected 20 kb product.	
Endonuclease Activity (Nicking, Polymerase, dNTP)	Pass



E0553S / Lot: 10086370

Page 1 of 2

Assay Name/Specification	Lot # 10086370
A 50 μl reaction in NEBuffer 2 in the presence of 200 μM dNTPs containing 1 μg of	
supercoiled PhiX174 DNA and a minimum of 10 units of Phusion® DNA Polymerase	
incubated for 4 hours at 37°C and 72°C results in <10% conversion to the nicked form	
as determined by agarose gel electrophoresis.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Christie Vazquez Production Scientist 12 Nov 2020

vistie Vazguez

Michael Tonello

Packaging Quality Control Inspector

12 Nov 2020