

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: SP6 RNA Polymerase

Catalog Number: M0207S
Concentration: 20,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to incorporate

1 nmol ATP into an acid-insoluble material in 1 hour at 37°C.

Packaging Lot Number: 10058293
Expiration Date: 08/2021
Storage Temperature: -20°C

Storage Conditions: 50 mM Tris-HCl, 100 mM NaCl, 20 mM \( \beta ME, 1 \) mM EDTA, 0.1 %

Triton®X-100, 50 % Glycerol, (pH 7.9 @ 25°C)

Specification Version: PS-M0207S/L v1.0

SP6 RNA Polymerase Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
M0207SVIAL	SP6 RNA Polymerase	10052056	Pass	
B9012SVIAL	RNAPol Reaction Buffer	10051065	Pass	

Assay Name/Specification	Lot # 10058293
RNase Activity (Extended Digestion) A 10 µl reaction in RNAPol Reaction Buffer containing 40 ng of a 300 base single-stranded RNA and a minimum of 20 units of SP6 RNA Polymerase is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass
Protein Purity Assay (SDS-PAGE) SP6 RNA Polymerase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
Promoter Specificity A 50 µl reaction in RNAPol Reaction Buffer in the presence of 2 mM NTPs containing 1 µg of Lambda DNA as a template and a minimum of 100 units of SP6 RNA Polymerase incubated for 1 hour at 37°C results in <1.5% of the amount of product incorporated as compared to a control reaction using SP6 DNA as a template.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in RNAPol Reaction Buffer containing 1 µg of Lambda DNA and a minimum of 100 units of SP6 RNA Polymerase incubated for 16 hours at 37°C results in	Pass



M0207S / Lot: 10058293

Page 1 of 2

Assay Name/Specification	Lot # 10058293
a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in RNAPol Reaction Buffer containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 100 units of SP6 RNA Polymerase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Endonuclease Activity (Nicking) A 50 µl reaction in RNAPol Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 100 units of SP6 RNA Polymerase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass

This product has been tested and shown to be in compliance with all specifications.

Dongxian Yue Production Scientist

13 Aug 2019

Josh Hersey

Packaging Quality Control Inspector

05 Dec 2019



M0207S / Lot: 10058293

Page 2 of 2