

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Mspl Methyltransferase

Catalog Number: M0215S
Concentration: 5,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme required to protect 1 µg

Lambda DNA in 1 hour at 37°C in a total reaction volume of 10 μl

against cleavage by Mspl restriction endonuclease.

Packaging Lot Number: 10208817
Expiration Date: 07/2025
Storage Temperature: -20°C

Storage Conditions: 50 mM Tris-HCl, 50 mM NaCl, 10 mM EDTA, 5 mM \( \beta ME, 200 \) \( \mu g/ml \) BSA, 50%

Glycerol, (pH 7.5 @ 25°C)

Specification Version: PS-M0215S v1.0

| Mspl Methyltransferase Component List |                                |            |                      |  |
|---------------------------------------|--------------------------------|------------|----------------------|--|
| <b>NEB Part Number</b>                | Component Description          | Lot Number | Individual QC Result |  |
| M0215SVIAL                            | Mspl Methyltransferase         | 10200156   | Pass                 |  |
| B9003SVIAL                            | S-adenosylmethionine (SAM)     | 10193027   | Pass                 |  |
| B0215SVIAL                            | Mspl Methylase Reaction Buffer | 10192345   | Pass                 |  |

| Assay Name/Specification  | Lot # 10208817 |
|---|----------------|
| Exonuclease Activity (Radioactivity Release) A 50 µl reaction in NEBuffer 2 containing 1 µg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 15 units of Mspl Methyltransferase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.   | Pass           |
| Functional Testing (Methyltransferase) A 10 μl reaction in Mspl Methyltransferase Reaction Buffer supplemented with 80 μM SAM containing 1 μg of Lambda DNA and 1 unit of Mspl Methyltransferase incubated for 1 hour at 37°C followed by heat inactivation results in ≥ 95% protection from digestion with 5 units of Mspl in NEBuffer 2 with 10 mM MgCl2 incubated at 37°C for 30 minutes as determined by agarose gel electrophoresis. | Pass           |
| Non-Specific DNase Activity (16 Hour) A 50 μl reaction in NEBuffer 2 containing 1 μg of Lambda-HindIII DNA and a minimum of 15 units of Mspl Methyltransferase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel   | Pass           |



M0215S / Lot: 10208817

Page 1 of 2



| Assay Name/Specification | Lot # 10208817 |
|--------------------------|----------------|
| electrophoresis.         |                |

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Jamie Souza

Production Scientist

2**5 √**ul 2023

Michael Tonello

Packaging Quality Control Inspector

28 Sep 2023

