240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: GpC Methyltransferase (M.CviPI)

Catalog #: M0227S/L
Concentration: 4,000 units/ml

Unit Definition: One unit is defined as the amount of enzyme required to protect 1 µg Lambda DNA in 1 hour at 37°C in a total reaction

volume of 10 µl against cleavage by HaeIII restriction endonuclease.

 Lot #:
 0041804

 Assay Date:
 04/2018

 Expiration Date:
 04/2020

 Storage Temp:
 -20°C

Storage Conditions: 15 mM Tris-HCl, 200 mM NaCl, 1 mM DTT, 0.1 mM EDTA, 50 % Glycerol, 200 µg/ml BSA, (pH 7.4 @ 25°C)

Specification Version: PS-M0227S/L v1.0
Effective Date: 21 May 2018

| Assay Name/Specification (minimum release criteria) | Lot #0041804 |
|---|--------------|
| Endonuclease Activity (Nicking) - A 50 μl reaction in GC Reaction Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 40 units of GpC Methyltransferase (M.CviPI) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Exonuclease Activity (Radioactivity Release) - A 50 μl reaction in GC Reaction Buffer containing 1 μg of a mixture of single and double-stranded [³ H] <i>E. coli</i> DNA and a minimum of 40 units of GpC Methyltransferase (M.CviPI) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Functional Testing (Methyltransferase) - A 20 μ l reaction in GC Reaction Buffer supplemented with 160 μ M SAM containing 1 μ g of Lambda DNA and 1 unit of GpC Methyltransferase (M.CviPI) incubated for 1 hour at 37°C followed by heat inactivation results in \geq 95% protection from digestion with 10 units of HaeIII in NEBuffer 2 incubated at 37°C for 1 hour as determined by agarose gel electrophoresis. | Pass |
| Non-Specific DNase Activity (16 Hour) - A 50 µl reaction in GC Reaction Buffer containing 1 µg of Lambda DNA and a minimum of 40 units of GpC Methyltransferase (M.CviPI) incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |

Authorized by Derek Robinson 21 May 2018







Inspected by Timothy Meixsell 03 Apr 2018

Timothy Meupelf