

New England Biolabs Certificate of Analysis

Product Name: Fpg
Catalog Number: M0240S
Concentration: 8,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to cleave 10 pmol of a 34-mer oligonucleotide duplex containing a single 8-oxoguanine base paired with a cytosine in a total reaction volume of 10 μ l in 1 hour at 37°C.
Lot Number: 10021307
Expiration Date: 09/2020
Storage Temperature: -20°C
Storage Conditions: 50 mM NaCl, 20 mM Tris-HCl, 0.5 mM EDTA, 50 % Glycerol, 200 μ g/ml BSA, (pH 8.0 @ 25°C)
Specification Version: PS-M0240S/L v1.0

| Fpg Component List | | | |
|--------------------|-----------------------|------------|----------------------|
| NEB Part Number | Component Description | Lot Number | Individual QC Result |
| M0240SVIAL | Fpg | 10020973 | Pass |
| B9001SVIAL | Purified BSA | 10014762 | Pass |
| B7001SVIAL | NEBuffer™ 1 | 0101804 | Pass |


| Assay Name/Specification | Lot # 10021307 |
|---|----------------|
| Exonuclease Activity (Radioactivity Release) A 50 μ l reaction in NEBuffer 1 containing 1 μ g of a mixture of single and double-stranded [³ H] E. coli DNA and a minimum of 24 units of Fpg incubated for 4 hours at 37°C releases <1.0% of the total radioactivity. | Pass |
| Non-Specific DNase Activity (16 Hour) A 50 μ l reaction in NEBuffer 1 containing 1 μ g of Lambda-HindIII DNA and a minimum of 40 units of Fpg incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis. | Pass |
| Protein Concentration (A280) The concentration of Fpg is 0.39 mg/ml +/- 10% as determined by UV absorption at 280 nm. Protein concentration is determined by the Pace method using the extinction coefficient of 39,795 and molecular weight of 30,290 daltons for Fpg (Pace, C.N. et al. (1995) Protein Sci., 4, 2411-2423). | Pass |

| Assay Name/Specification | Lot # 10021307 |
|--|----------------|
| <p>Protein Purity Assay (SDS-PAGE) Fpg is \geq 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p> | Pass |
| <p>qPCR DNA Contamination (E. coli Genomic) A minimum of 8 units of Fpg is screened for the presence of E. coli genomic DNA using SYBR[®] Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is \leq 1 E. coli genome.</p> | Pass |
| <p>RNase Activity (Extended Digestion) A 10 μl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 8 units of Fpg is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p> | Pass |

This product has been tested and shown to be in compliance with all specifications.



Tony Spear-Alfonso
Production Scientist
12 Sep 2018



Michael Tonello
Packaging Quality Control Inspector
02 Oct 2018