

New England Biolabs Certificate of Analysis


Product Name: *Therminator™ DNA Polymerase*
Catalog Number: *M0261S*
Concentration: *2,000 U/ml*
Unit Definition: *One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid insoluble material in 30 minutes at 75°C.*
Lot Number: *10042500*
Expiration Date: *04/2021*
Storage Temperature: *-20°C*
Storage Conditions: *10 mM Tris-HCl , 100 mM KCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol, (pH 7.4 @ 25°C)*
Specification Version: *PS-M0261S/L v1.0*

Therminator™ DNA Polymerase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0261SVIAL	Therminator™ DNA Polymerase	10042499	Pass
B9004SVIAL	ThermoPol® Reaction Buffer Pack	0031712	Pass

Assay Name/Specification	Lot # 10042500
<p>Endonuclease Activity (Nicking) A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 units of Therminator™ DNA Polymerase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 20 units of Therminator™ DNA Polymerase incubated for 4 hours at 37°C and 75°C releases <0.1% of the total radioactivity.</p>	Pass
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 2 units of Therminator™ DNA Polymerase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>Phosphatase Activity (pNPP)</p>	Pass

Assay Name/Specification	Lot # 10042500
<p>A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl₂ containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units Therminator™ DNA Polymerase incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p>	
<p>Protein Purity Assay (SDS-PAGE) Therminator™ DNA Polymerase is ≥ 98% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Therminator™ DNA Polymerase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.



Doreen Duquette
Production Scientist
19 Apr 2019



Josh Hersey
Packaging Quality Control Inspector
13 May 2019