

## New England Biolabs Certificate of Analysis

**Product Name:** *Therminator™ DNA Polymerase*  
**Catalog Number:** *M0261S*  
**Concentration:** *2,000 U/ml*  
**Unit Definition:** *One unit is defined as the amount of enzyme that will incorporate 10 nmol of dNTP into acid insoluble material in 30 minutes at 75°C.*  
**Packaging Lot Number:** *10200477*  
**Expiration Date:** *07/2025*  
**Storage Temperature:** *-20°C*  
**Storage Conditions:** *10 mM Tris-HCl , 100 mM KCl , 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol, (pH 7.4 @ 25°C)*  
**Specification Version:** *PS-M0261S/L v2.0*


Therminator™ DNA Polymerase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0261SVIAL	Therminator™ DNA Polymerase	10198991	Pass
B9004SVIAL	ThermoPol® Reaction Buffer Pack	10187437	Pass

Assay Name/Specification	Lot # 10200477
<p><b>Endonuclease Activity (Nicking)</b>            A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 units of Therminator™ DNA Polymerase incubated for 4 hours at 37°C results in &lt;10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p><b>Exonuclease Activity (Radioactivity Release)</b>            A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of a mixture of single and double-stranded [<sup>3</sup>H] E. coli DNA and a minimum of 20 units of Therminator™ DNA Polymerase incubated for 4 hours at 37°C and 75°C releases &lt;0.1% of the total radioactivity.</p>	Pass
<p><b>Non-Specific DNase Activity (16 Hour)</b>            A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 2 units of Therminator™ DNA Polymerase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p><b>Phosphatase Activity (pNPP)</b></p>	Pass

Assay Name/Specification	Lot # 10200477
<p>A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl<sub>2</sub> containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units Therminator™ DNA Polymerase incubated for 4 hours at 37°C yields &lt;0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p>	
<p><b>Protein Purity Assay (SDS-PAGE)</b> Therminator™ DNA Polymerase is ≥ 98% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	<b>Pass</b>
<p><b>RNase Activity (Extended Digestion)</b> A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Therminator™ DNA Polymerase is incubated at 37°C. After incubation for 16 hours, &gt;90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	<b>Pass</b>

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit [www.neb.com/trademarks](http://www.neb.com/trademarks) for additional information.

  
 Lea Antonopoulos  
 Production Scientist  
 29 Jul 2023

  
 Josh Hersey  
 Packaging Quality Control Inspector  
 31 Jul 2023