240 County Road
Ipswich, MA 01938-2723
Tel 978-927-5054
Fax 978-921-1350
www.neb.com
info@neb.com

New England Biolabs
Certificate of Analysis

Product Name:	*Taq DNA Polymerase with Standard Taq Buffer*
Catalog Number:	*M0273E*
Concentration:	*5,000 U/ml*
Unit Definition:	*One unit is defined as the amount of enzyme that will incorporate 15 nmol of dNTP into acid insoluble material in 30 minutes at 75°C.*
Packaging Lot Number:	*10269419*
Expiration Date:	*11/2026*
Storage Temperature:	*-20°C*
Storage Conditions:	*10 mM Tris-HCl , 100 mM KCl , 1 mM DTT , 0.1 mM EDTA , 0.5 % Tween® 20 , 0.5 % IGEPAL® CA-630 , 50 % Glycerol, (pH 7.4 @ 25°C)*
Specification Version:	*PS-M0273S/L/X/E/G/V v3.0*

Taq DNA Polymerase with Standard Taq Buffer Component List			
NEB Part Number	**Component Description**	**Lot Number**	**Individual QC Result**
M0273XVIAL	Taq DNA Polymerase with Standard Taq Buffer	10264523	**Pass**
B9014SVIAL	Standard Taq Reaction Buffer Pack	10246931	**Pass**

Assay Name/Specification	**Lot # 10269419**
Endonuclease Activity (Nicking) A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 units of Taq DNA Polymerase incubated for 4 hours at 37°C and 75ºC results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	**Pass**
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 5 units of Taq DNA Polymerase incubated for 16 hours at 37ºC results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.	**Pass**
PCR Amplification (5.0 kb Lambda DNA) A 50 µl reaction in Standard Taq Reaction Buffer in the presence of 200 µM dNTPs and 0.2 µM primers containing 5 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 5.0 kb product.	**Pass**
Phosphatase Activity (pNPP) A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl2 containing 2.5 mM	**Pass**

NEW ENGLAND Biolabs®

240 County Road
Ipswich, MA 01938-2723

Tel 978-927-5054
Fax 978-921-1350

www.neb.com
info@neb.com

Assay Name/Specification	Lot # 10269419
p-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units Taq DNA Polymerase incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	
Protein Purity (Microfluidic Electrophoresis) Taq DNA Polymerase is ≥97% pure as determined by microfluidic electrophoresis.	**Pass**
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Taq DNA Polymerase is incubated at 37ºC. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	**Pass**
Single Stranded DNase Activity (FAM-Labeled Oligo) A 50 µl reaction in ThermoPol® Reaction Buffer containing a 10 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 25 units of Taq DNA Polymerase incubated for 30 minutes at 37°C and 75°C yields <10% degradation as determined by capillary electrophoresis.	**Pass**
qPCR DNA Contamination (E. coli Genomic) A minimum of 5 units of Taq DNA Polymerase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	**Pass**

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Trinh Nguyen
Production Scientist
31 Dec 2024

Michael Tonello
Packaging Quality Control Inspector
17 Jan 2025