

New England Biolabs Certificate of Analysis

Product Name: AMV Reverse Transcriptase
Catalog Number: M0277S
Concentration: 10,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme required to incorporate 1 nmol of dTTP into an acid-insoluble form in 10 minutes at 37°C.
Packaging Lot Number: 10208720
Expiration Date: 10/2025
Storage Temperature: -20°C
Storage Conditions: 200 mM KPO₄, 2 mM DTT, 0.2 % Triton®X-100, 50 % Glycerol, (pH 7.2 @ 25°C)
Specification Version: PS-M0277S/L v2.0


AMV Reverse Transcriptase Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0277SVIAL	AMV Reverse Transcriptase	10203129	Pass
B0277AVIAL	AMV Reverse Transcriptase Reaction Buffer	10177226	Pass

Assay Name/Specification	Lot # 10208720
<p>Endonuclease Activity (Nicking) A 50 µl reaction in AMV Reverse Transcriptase Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 50 units of AMV Reverse Transcriptase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>Exonuclease Activity (Radioactivity Release) A 50 µl reaction in AMV Reverse Transcriptase Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of AMV Reverse Transcriptase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.</p>	Pass
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 or T7 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 10 units of AMV Reverse Transcriptase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>RNase Activity Assay (4 Hour Digestion)</p>	Pass

Assay Name/Specification	Lot # 10208720
A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of AMV Reverse Transcriptase is incubated at 37°C. After incubation for 4 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.



Lea Antonopoulos
Production Scientist
01 Nov 2023



Michael Tonello
Packaging Quality Control Inspector
02 Nov 2023