

New England Biolabs Certificate of Analysis

Product Name: *Afu Uracil-DNA Glycosylase (UDG)*
Catalog #: *M0279S/L*
Concentration: *2,000 units/ml*
Unit Definition: *One unit is defined as the amount of enzyme that catalyzes the release of 60 pmol of uracil per minute from double-stranded, uracil-containing DNA. Activity is measured by release of [³H]-uracil in a 50 µl reaction containing 0.2 µg DNA (10⁴-10⁵ cpm/µg) in 30 minutes at 65°C.*
Lot #: *0031501*
Assay Date: *01/2015*
Expiration Date: *1/2017*
Storage Temp: *Store between -25 and -15 °C*
Storage Conditions: *10 mM Tris-HCl, 50 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.1 mg/ml BSA, 50% Glycerol, pH 7.4 @ 25°C*
Specification Version: *PS-M0279S/L v1.0*
Effective Date: *14 Feb 2014*

Assay Name/Specification (minimum release criteria)	Lot #0031501
DNase Activity (Labeled Oligo, 3' extension) - A 50 µl reaction in CutSmart™ Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 3' extension and a minimum of 10 units of Afu Uracil-DNA Glycosylase (UDG) incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	Pass
DNase Activity (Labeled Oligo, 5' extension) - A 50 µl reaction in CutSmart™ Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a 5' extension and a minimum of 10 units of Afu Uracil-DNA Glycosylase (UDG) incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	Pass
Double Stranded DNase Activity (Labeled Oligo) - A 50 µl reaction in CutSmart™ Buffer containing a 20 nM solution of a fluorescent labeled double-stranded oligonucleotide containing a blunt end and a minimum of 10 units of Afu Uracil-DNA Glycosylase (UDG) incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.	Pass
Endonuclease Activity (Nicking) - A 50 ul reaction in ThermoPol Reaction Buffer containing 1 ug of supercoiled PhiX174 DNA and a minimum of 4 units of Afu Uracil-DNA Glycosylase (UDG) incubated for 4 hours at 65°C results in <20% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) - A 50 µl reaction in ThermoPol Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³ H] <i>E. coli</i> DNA and a minimum of 50 units of Afu Uracil-DNA Glycosylase (UDG) incubated for 4 hours at 65°C releases <0.1% of the total radioactivity.	Pass



New England Biolabs Certificate of Analysis

Assay Name/Specification (minimum release criteria)	Lot #0031501
<p>Non-Specific DNase Activity (16 Hour) - A 50 μl reaction in ThermoPol Reaction Buffer containing 1 μg of Lambda DNA and a minimum of 50 units of Afu Uracil-DNA Glycosylase (UDG) incubated for 16 hours at 65°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>Protein Concentration (A280/NanoDrop) - The concentration of Afu Uracil-DNA Glycosylase (UDG) is 0.027 mg/ml +/- 10% as determined by UV absorption at 280 nm. Protein concentration is determined by the Pace method using the extinction coefficient of 16,110 and molecular weight of 22,720 daltons for Afu Uracil-DNA Glycosylase (UDG) (Pace, C.N. et al. (1995) Protein Sci., 4, 2411-2423).</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) - Afu Uracil-DNA Glycosylase (UDG) is \geq 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>RNase Activity (Extended Digestion) - A 10 μl reaction in NEBuffer 4 containing 40 ng of a 300 base single stranded RNA and a minimum of 2 units of Afu Uracil-DNA Glycosylase (UDG) is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using polyacrylamide gel electrophoresis detection.</p>	Pass
<p>Single Stranded DNase Activity (FAM-Labeled Oligo) - A 50 μl reaction in CutSmart™ Buffer containing a 20 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 10 units of Afu Uracil-DNA Glycosylase (UDG) incubated for 16 hours at 37°C yields <5% degradation as determined by capillary electrophoresis.</p>	Pass

* The BSA in this product has been granted an EDQM "Certificate of Suitability" from the European Directorate for the Quality of Medicines (# R1-CEP-2003-204-Rev00) and has been granted a USDA Certificate for Export of Bovine Blood Plasma/Serum for Manufacture into Pharmaceutical Products.



Authorized by
Derek Robinson
14 Feb 2014



Inspected by
Lauren Sears
05 Feb 2015

