

be INSPIRED

ve DISCOVERY 240 County Road

ve GENUINE Inswich MA 01938

240 County Road Tel 978-927-5054 Ipswich, MA 01938-2723 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Antarctic Phosphatase

Catalog Number: M0289L Concentration: 5,000 U/ml

Unit Definition: One unit is defined as the amount of enzyme that will

dephosphorylate 1 µg of pUC19 vector DNA cut with a restriction enzyme generating 5´ recessed ends in 30 minutes at 37°C. Dephosphorylation is defined as >95% inhibition of recircularization

in a self-ligation reaction and is measured by transfor

Lot Number: 10014915
Expiration Date: 06/2020
Storage Temperature: -20°C

Storage Conditions: 10 mM Tris-HCl, 1 mM MgCl2, 0.01 mM ZnCl2, 50 % Glycerol, (pH 7.4 @

25°C)

Specification Version: PS-M0289S/L v1.0

Antarctic Phosphatase Component List				
NEB Part Number	Component Description	Lot Number	Individual QC Result	
M0289LVIAL	Antarctic Phosphatase	10009601	Pass	
B0289SVIAL	Antarctic Phosphatase Reaction Buffer	0101709	Pass	

Assay Name/Specification	Lot # 10014915
Endonuclease Activity (Nicking) A 50 μl reaction in Antarctic Phosphatase Reaction Buffer containing 1 μg of supercoiled PhiX174 DNA and a minimum of 50 units of Antarctic Phosphatase incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 µl reaction in Antarctic Phosphatase Reaction Buffer containing 1 µg of a mixture of single and double-stranded [³H] E. coli DNA and a minimum of 50 units of Antarctic Phosphatase incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 4 containing 1 µg of PhiX174-HaeIII DNA and a minimum of 50 units of Antarctic Phosphatase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel	Pass



M0289L / Lot: 10014915

Page 1 of 2

Assay Name/Specification	Lot # 10014915
electrophoresis.	
Protein Purity Assay (SDS-PAGE) Antarctic Phosphatase is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	Pass
qPCR DNA Contamination (E. coli Genomic) A minimum of 5 units of Antarctic Phosphatase is screened for the presence of E. coli genomic DNA using SYBR® Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.	Pass
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Antarctic Phosphatase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

Heidi Church

Production Scientist

10 Jul 2018

Michael Tonello

Packaging Quality Control Inspector

16 Jul 2018



M0289L / Lot: 10014915

Page 2 of 2