

New England Biolabs Certificate of Analysis

Product Name: Taq DNA Polymerase with Standard Taq (Mg-free) Buffer
Catalog Number: M0320L
Concentration: 5,000 U/ml
Unit Definition: One unit is defined as the amount of enzyme that will incorporate 15 nmol of dNTP into acid-insoluble material in 30 minutes at 75°C.
Lot Number: 10013987
Expiration Date: 06/2020
Storage Temperature: -20°C
Storage Conditions: 10 mM Tris-HCl, 100 mM KCl, 1 mM DTT, 0.1 mM EDTA, 0.5 % Tween® 20, 0.5 % IGEPAL® CA-630, 50 % Glycerol, (pH 7.4 @ 25°C)
Specification Version: PS-M0320S/L v1.0

Taq DNA Polymerase with Standard Taq (Mg-free) Buffer Component List			
NEB Part Number	Component Description	Lot Number	Individual QC Result
M0320LVIAL	Taq DNA Polymerase with Standard Taq (Mg-free) Buffer	10009973	Pass
B9021SVIAL	Magnesium Chloride (MgCl ₂) Solution	0031705	Pass
B9015SVIAL	Standard Taq (Mg-free) Reaction Buffer Pack	0021703	Pass


Assay Name/Specification	Lot # 10013987
<p>Endonuclease Activity (Nicking) A 50 µl reaction in ThermoPol® Reaction Buffer containing 1 µg of supercoiled PhiX174 DNA and a minimum of 20 units of Taq DNA Polymerase incubated for 4 hours at either 37°C or 75°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.</p>	Pass
<p>Non-Specific DNase Activity (16 Hour) A 50 µl reaction in NEBuffer 2 containing 1 µg of T3 DNA in addition to a reaction containing Lambda-HindIII DNA and a minimum of 5 units of Taq DNA Polymerase incubated for 16 hours at 37°C results in a DNA pattern free of detectable nuclease degradation as determined by agarose gel electrophoresis.</p>	Pass
<p>PCR Amplification (5.0 kb Lambda DNA) A 50 µl reaction in Standard Taq (Mg-free) Reaction Buffer plus 1.5 mM MgCl₂ in the presence of 200 µM dNTPs and 0.2 µM primers containing 5 ng Lambda DNA with 1.25 units of Taq DNA Polymerase for 25 cycles of PCR amplification results in the expected 5.0 kb product.</p>	Pass

Assay Name/Specification	Lot # 10013987
<p>Phosphatase Activity (pNPP) A 200 µl reaction in 1M Diethanolamine, pH 9.8, 0.5 mM MgCl₂ containing 2.5 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 100 units Taq DNA Polymerase incubated for 4 hours at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.</p>	Pass
<p>Protein Purity Assay (SDS-PAGE) Taq DNA Polymerase is ≥ 99% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.</p>	Pass
<p>qPCR DNA Contamination (E. coli Genomic) A minimum of 5 units of Taq DNA Polymerase is screened for the presence of E. coli genomic DNA using SYBR[®] Green qPCR with primers specific for the E. coli 16S rRNA locus. Results are quantified using a standard curve generated from purified E. coli genomic DNA. The measured level of E. coli genomic DNA contamination is ≤ 1 E. coli genome.</p>	Pass
<p>RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 1 µl of Taq DNA Polymerase is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.</p>	Pass
<p>Single Stranded DNase Activity (FAM-Labeled Oligo) A 50 µl reaction in ThermoPol[®] Reaction Buffer containing a 10 nM solution of a fluorescent internal labeled oligonucleotide and a minimum of 25 units of Taq DNA Polymerase incubated for 30 minutes at either 37°C or 75°C yields <10% degradation as determined by capillary electrophoresis.</p>	Pass

This product has been tested and shown to be in compliance with all specifications.



Lynne Apone
Production Scientist
21 Jun 2018



Michael Tonello
Packaging Quality Control Inspector
26 Jul 2018