

240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

## New England Biolabs Certificate of Analysis

Product Name: Pyrophosphatase, Inorganic (E. coli)

Catalog Number: M0361L Concentration: 100 U/ml

Unit Definition: One unit is the amount of enzyme that will generate 1 µmol of

phosphate per minute from inorganic pyrophosphate under standard

reaction conditions.

Packaging Lot Number: 10092476
Expiration Date: 10/2022
Storage Temperature: -20°C

Storage Conditions: 100 mM NaCl , 20 mM Tris-HCl (pH 8.0), 1 mM DTT , 0.1 mM EDTA , 50 %

Glycerol

Specification Version: PS-M0361S/L v2.0

Pyrophosphatase, Inorganic (E. coli) Component List				
<b>NEB Part Number</b>	Component Description	Lot Number	Individual QC Result	
M0361LVIAL	Pyrophosphatase, Inorganic (E. coli)	10086151	Pass	

Assay Name/Specification	Lot # 10092476
Endonuclease Activity (Nicking) A 50 μl reaction in NEBuffer 4 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 0.5 units of Pyrophosphatase, Inorganic (E. coli) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis.	Pass
Exonuclease Activity (Radioactivity Release) A 50 μl reaction in NEBuffer 4 containing 1 μg of a mixture of single and double-stranded [ ³H] E. coli DNA and a minimum of 0.5 units of Pyrophosphatase, Inorganic (E. coli) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity.	Pass
Phosphatase Activity (pNPP) A 100 µl reaction in NEBuffer 4 containing 10 mM p-Nitrophenyl Phosphate (pNPP) and a minimum of 1 unit Pyrophosphatase, Inorganic (E. coli) incubated for 1 hour at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis.	Pass
Protein Purity Assay (SDS-PAGE)	Pass



M0361L / Lot: 10092476

Page 1 of 2

Assay Name/Specification	Lot # 10092476
Pyrophosphatase, Inorganic (E. coli) is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection.	
RNase Activity (Extended Digestion) A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 0.1 unit of Pyrophosphatase, Inorganic (E. coli) is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection.	Pass

This product has been tested and shown to be in compliance with all specifications.

One or more products referenced in this document may be covered by a 3rd-party trademark. Please visit www.neb.com/trademarks for additional information.

Bhairavi Jani Production Scientist 15 Dec 2020 Michael Tonello

Packaging Quality Control Inspector

15 Dec 2020



M0361L / Lot: 10092476

Page 2 of 2