240 County Road Ipswich, MA 01938-2723 Tel 978-927-5054 Fax 978-921-1350 www.neb.com info@neb.com

New England Biolabs Certificate of Analysis

Product Name: Pyrophosphatase, Inorganic (E. coli)

Catalog #: M0361S/L
Concentration: 100 units/ml

Unit Definition: One unit is the amount of enzyme that will generate 1 µmol of phosphate per minute from inorganic pyrophosphate under

standard reaction conditions.

 Lot #:
 0031803

 Assay Date:
 03/2018

 Expiration Date:
 03/2020

 Storage Temp:
 -20°C

Storage Conditions: 100 mM NaCl , 20 mM Tris-HCl (pH 8.0), 1 mM DTT , 0.1 mM EDTA , 50 % Glycerol

Specification Version: PS-M0361S/L v2.0
Effective Date: 27 Apr 2015

| Assay Name/Specification (minimum release criteria) | Lot #0031803 |
|--|--------------|
| Endonuclease Activity (Nicking) - A 50 μl reaction in NEBuffer 4 containing 1 μg of supercoiled PhiX174 DNA and a minimum of 0.5 units of Pyrophosphatase, Inorganic (<i>E. coli</i>) incubated for 4 hours at 37°C results in <10% conversion to the nicked form as determined by agarose gel electrophoresis. | Pass |
| Exonuclease Activity (Radioactivity Release) - A 50 μl reaction in NEBuffer 4 containing 1 μg of a mixture of single and double-stranded [³ H] <i>E. coli</i> DNA and a minimum of 0.5 units of Pyrophosphatase, Inorganic (<i>E. coli</i>) incubated for 4 hours at 37°C releases <0.1% of the total radioactivity. | Pass |
| Phosphatase Activity (pNPP) - A 100 μl reaction in NEBuffer 4 containing 10 mM <i>p</i> -Nitrophenol Phosphate (pNPP) and a minimum of 1 unit Pyrophosphatase, Inorganic (<i>E. coli</i>) incubated for 1 hour at 37°C yields <0.0001 unit of alkaline phosphatase activity as determined by spectrophotometric analysis. | Pass |
| Protein Purity Assay (SDS-PAGE) - Pyrophosphatase, Inorganic (<i>E. coli</i>) is ≥ 95% pure as determined by SDS-PAGE analysis using Coomassie Blue detection. | Pass |
| RNase Activity (Extended Digestion) - A 10 µl reaction in NEBuffer 4 containing 40 ng of a 300 base single-stranded RNA and a minimum of 0.1 unit of Pyrophosphatase, Inorganic (<i>E. coli</i>) is incubated at 37°C. After incubation for 16 hours, >90% of the substrate RNA remains intact as determined by gel electrophoresis using fluorescent detection. | Pass |

Authorized by Derek Robinson 27 Apr 2015







Inspected by
Dongxian Yue
21 Mar 2018